


CareerGuide.com

BITSAT 2023

Sample Paper

Physics

- Q.1 The Figure represents a voltage regulator circuit using of the Zener diode is $6V$ and the load resistance is $R_L = 4K\Omega$. The series resistance $R_i = 1K\Omega$. If the battery V_B varies from $8V$ to $16V$, What are the minimum and maximum values of the current respective



Option 1:

$0.5mA; 6mA$

Option 2:

$1mA; 8.5mA$

Option 3:

$0.5mA; 8.5mA$

Option 4:

$1.5mA; 8.5mA$

Correct Answer:

$0.5mA; 8.5mA$

Solution:

Given $V_B = 8V$

$$i_L = \frac{6 \times 10^{-3}}{4} = 1.5 \times 10^{-3} A \text{ and } i_R = \frac{(8 - 6) \times 10^{-3}}{1} = 2 \times 10^{-3} A$$

$$\therefore i_{\text{Zener diode}} = i_R - i_L = 0.5 \times 10^{-3} A$$

When $V_B = 16V$

$$i_L = 1.5 \times 10^{-3} A \text{ and } i_R = \frac{(16 - 6) \times 10^{-3}}{1} = 10 \times 10^{-3} A$$

$$\therefore i_{\text{Zener diode}} = i_R - i_L = 8.5 \times 10^{-3} A$$

Q.2

10 gm of ice cubes at 0°C are released in a tumbler (water at 55°C). Assuming that negligible heat is taken from the surroundings, the temperature of the tumbler becomes nearly

Option 1:

31°C

Option 2:

22°C

Option 3:

19°C

Option 4:

15°C

Correct Answer:

22°C

Solution:

As we have learned

Water Equivalent -

$w = mc$ gm

- wherein

$C_{\text{water}} = 1$

Heat gained = heat lost

heat gained $(\theta - 0)L + ms$ (

$= 10 \times 80 + 10 \times 1 \times \theta$

Heat lost by water = $55(40 - \theta)$

$\Rightarrow 800 + 10\theta = 2200 - 550 \text{ or } 650 = 1400$

$\theta = 22^\circ\text{C}$

Q.3

If the magnetic dipole moment of an atom of the diamagnetic material, and ferromagnetic material is predicted by

Option 1:

$$\mu_d \neq 0 \text{ and } \mu_f \neq 0$$

Option 2:

$$\mu_p = 0 \text{ and } \mu_f \neq 0$$

Option 3:

$$\mu_d = 0 \text{ and } \mu_p \neq 0$$

Option 4:

$$\mu_d \neq 0 \text{ and } \mu_p = 0$$

Correct Answer:

$$\mu_d = 0 \text{ and } \mu_p \neq 0$$

Solution:

As we learned

Magnetic Permeability -

It is the degree or extent to which a magnetic field is induced. It is

-

Properties of substances

Q.4

Distance of the centre of mass of a solid uniform cone of its base is R and its height is h then z_0 is equal to :

Option 1:

$$\frac{h^2}{4R}$$

Option 2:

$$\frac{3h}{4}$$

Option 3:

$$\frac{5h}{8}$$

Option 4:

$$\frac{3h^2}{8R}$$


Correct Answer:

$$\frac{3h}{4}$$

Solution:

As we learnt in

Centre of Mass of a solid cone from the vertex is $\frac{3h}{4}$.



Q. 5 Two plates are 2 cm apart, a potential difference of 10 volt is applied between them, the electric field between the plates is

Option 1:

$$20\text{ N/C}$$

Option 2:

$$500\text{ N/C}$$

Option 3:

$$5\text{ N/C}$$

Option 4:

250 N/C

Correct Answer:

500 N/C


Solution:

As we learned

Electric Field Intensity -

$$\vec{E} = \frac{\vec{F}}{q_0} = \frac{kQ}{r^2}$$

- wherein



$$E = \frac{V}{d} = \frac{10}{2 \times 10^{-2}} = 500 \text{ N/C}$$

Q.6

An electric field $\vec{E} = (25\hat{i} + 30\hat{j}) \text{ NC}^{-1}$ exists in a region of space. If the origin is taken to be zero then the potential at $x=2$

Option 1:

-130 J

Option 2:

-120 J

Option 3:

- 140 J

Option 4:

- 110 J

Correct Answer:

- 110 J

Solution:

As we learnt in

In space -

$$E_x = \frac{-dv}{dx} \quad E_y = \frac{-dv}{dy} \quad , \quad E_z = \frac{-dv}{dz}$$

-

$$dv = -\vec{E} \cdot d\vec{r} \Rightarrow \int dV = - \int \vec{E} \cdot d\vec{r}$$

$$d\vec{r} = dx\hat{i} + dy\hat{j} + dz\hat{k}$$

$$\vec{E} = (25\hat{i} + 30\hat{j})Nc^{-1}$$

$$\int dV = - \int (25\hat{i} + 30\hat{j})(dx\hat{i} + dy\hat{j})$$

$$\int_0^V dV = \int_0^2 25dx + \int_0^2 30dy$$


$$V - 0 = 25[x]_0^2 + 30[y]_0^2$$

$$V = -[25 \times 2 + 30 \times 2] \Rightarrow V = -110J / C$$

$$V = -110J / C$$

Q.7

Four capacitors of each of capacity $3\mu\text{F}$ are connected as shown in the figure. The ratio of equivalent capacitance between A and B and between A and C is



Option 1:

4 : 3

Option 2:

3 : 4

Option 3:

2 : 3

Option 4:

3 : 2

Correct Answer:

3 : 4


Solution:

As we learnt in

Series Grouping -

$$\frac{1}{C_{eq}} = \frac{1}{C_1} + \frac{1}{C_2} + \dots$$

- wherein




and

Parallel Grouping -

$$C_{eq} = C_1 + C_2 + \dots$$

- wherein



Between A and B, 3 capacitors are in series and they are in p

$$C_1 = \frac{C \times 3C}{C + 3C} = \frac{3C}{4}$$


Between A and C:

Two capacitors are in series and in turn they are parallel.

$$C_2 = \frac{2C \times 2C}{2C + 2C} = C$$

$$\text{Ratio} = \frac{3}{4}$$

Q.8 The given graph shows variation (with distance r from c



Option 1:

potential of a uniformly charged spherical shell

Option 2:

Electric eld of a uniformly charged spherical shell

Option 3:

Electric field of a uniformly charged sphere

Option 4:

Potential of a uniformly charged sphere

Correct Answer:

potential of a uniformly charged spherical shell


Solution:

Inside the sphere (P lies inside the sphere) -

$$E_{in} = 0$$

$$V_{in} = \text{constant}$$

- wherein



The potential of uniformly charged spherical shell.

Q.9

A rod, of length L at room temperature and uniform area of cross-section A is expanded by an external compressive force F , is applied on each of its ends, preventing it from contracting. The temperature of the rod is raised by ΔT . The Young's modulus, Y , for this material is

Option 1:

$$\frac{F}{A\alpha\Delta T}$$

Option 2:

$$\frac{F}{A\alpha(\Delta T - 273)}$$

Option 3:

$$\frac{F}{2A\alpha\Delta T}$$

Option 4:

$$\frac{2F}{A\alpha\Delta T}$$

Correct Answer:

$$\frac{F}{A\alpha\Delta T}$$

Solution:

Elongation -

$$\Delta l = \frac{FL}{AY}$$

- wherein

$$\Delta l = \text{Elongation}$$

$$y = \frac{\frac{F}{A}}{\frac{\Delta L}{L}}$$

$$\Rightarrow \frac{\Delta L}{L} = \frac{\frac{F}{A}}{y}$$

Coefficient of linear expansion

$$\alpha = \frac{\Delta L}{L\Delta T}$$

$$\frac{\Delta L}{L} = \alpha\Delta T$$


So we get

$$y = \frac{F}{A\alpha\Delta T}$$


Q. 10

The form factor for which of these is $\frac{\pi}{2\sqrt{2}}$


Option 1:




Option 2:




Option 3:



Option 4:



Correct Answer:



Solution:


As we learnt

Wave Form (Sinusoidal) -

$$\text{R.M.S. Value} = \frac{i_0}{\sqrt{2}}$$

$$\text{average value} = \frac{2i_0}{\pi}$$

- wherein



$$\text{form factor} = \frac{\text{rms value}}{\text{average value}}$$

For sinusoidal wave

$$\text{R.M.S. Value} = \frac{i_0}{\sqrt{2}}$$

$$\text{average value} = \frac{2i_0}{\pi}$$

$$R_f = \frac{\frac{i_0}{\sqrt{2}}}{\frac{2i_0}{\pi}} \Rightarrow R_f = \frac{\pi}{2\sqrt{2}}$$

Q.11 An electron moving with speed parallel to the electric field of intensity 10^3 N/C . The field is responsible for the retardation of electrons. Now evaluate the distance traveled by the electron in 1 ns (mass of $e = 9.1 \times 10^{-31} \text{ kg}$ charge $e = 1.6 \times 10^{-19} \text{ C}$)

Option 1:

7 m

Option 2:

0.7 mm

Option 3:

7 cm

Option 4:

0.7 cm

Correct Answer:

7 cm

Solution:

As we have learnt,

when Charged Particle at rest in uniform eld -

Force and acceleration

$$F = QE$$

$$a = \frac{QE}{m}$$

- wherein

m - mass

Q - charge


E - Electric eld strength.

$$\text{Electric force} \Rightarrow a = \frac{QE}{m} \therefore a = \frac{1.6 \times 10^{-19} \times 1 \times 10^3}{9 \times 10^{-31}} = \frac{1.6}{9} \times 10^{15}$$

$$u = 5 \times 10^6 \text{ and } 0 \therefore \text{from } v^2 = u^2 - 2as \Rightarrow s = \frac{u^2}{2a}$$

$$\therefore \text{Distance } s = \frac{(5 \times 10^6)^2 \times 9}{2 \times 1.6 \times 10^{15}} = 7 \text{ cm (approx)}$$

Q.12 In the circuit shown in figure, find the current through the



Option 1:

5 A

Option 2:

0 A

Option 3:

3 A

Option 4:

4 A

Correct Answer:

5 A

Solution:


As we have learnt,

Kirchhoff's Law -

$$\sum q = 0$$

-

The current in the circuit are assumed as shown in the g.



Applying KVL along the loop $ABDA$, we get

$$-6i_1 - 3i_2 + 15 = 0 \text{ or } 2i_1 + i_2 = 5 \dots\dots(i)$$

Applying KVL along the loop $BCDB$, we get

$$-3(i1 - i2) - 30 + 3i2 = 0 \text{ or } -i1 + 2i2 = 10 \dots\dots(ii)$$

Solving equation (i) and (ii) for i_2 , we get $i_2 = 5 \text{ A}$.

Q.13 When a ball is thrown up vertically with velocity v_0 it reaches a maximum height h . If it wishes to triple the maximum height then the ball should be thrown with velocity

Option 1:

$$v \circ \sqrt{3}$$

Option 2:

3 v o

Option 3:

9 v o

Option 4:

$$(3/2) v_0$$

Correct Answer:

$v \circ \sqrt{3}$

Solution:

As we learnt in

Potential Energy -

$$U_f - U_i = \int_{r_i}^{r_f} \vec{f} \cdot d\vec{s}$$

- wherein

U_f – final potential energy

U_i – initial potential energy

f – force

ds – small displacement

r_i – initial position

r_f – final position

At Maximum height $V = 0$

$$v_0 = \sqrt{2gh} \text{ --- (i)}$$

$$\text{At } h' = 3h$$


$$u^2 = 2g(3h) = 6gh$$

$$u = \sqrt{6gh} \text{ --- (ii)}$$

$$\frac{v_0}{u} = \frac{\sqrt{2gh}}{\sqrt{6gh}}$$

$$\Rightarrow u = \sqrt{3}v_0$$

Q. 14 A rigid square of side a and carrying current I_2 is lying on a horizontal surface long carrying wire in the same plane as shown in figure. The force on the square due to the wire will be :



Option 1:

Repulsive and equal to $\frac{\mu_0 I_1 I_2}{2\pi}$

Option 2:

Attractive and equal to $\frac{\mu_0 I_1 I_2}{3\pi}$

Option 3:

Repulsive and equal to $\frac{\mu_0 I_1 I_2}{4\pi}$

Option 4:

Zero

Correct Answer:

Repulsive and equal to $\frac{\mu_0 I_1 I_2}{4\pi}$


Solution:

Force between two parallel current carrying conductors -

$$F = \frac{\mu}{4\pi} \frac{2I_1 I_2 l}{a}$$


$$\frac{F}{l} = \frac{\mu_0 2I_1 I_2}{4\pi a}$$

- wherein



I_1 and I_2 current carrying two parallel wires

a - separation between two wires



$$F_1 = \frac{\mu_0 I_1 I_2 a}{2\pi a} = \frac{\mu_0 I_1 I_2}{2\pi}$$

$$F_2 = \frac{\mu_0 I_1 I_2 a}{2\pi \cdot 2a} = \frac{\mu_0 I_1 I_2}{4\pi}$$

$$F_{Total} = F_1 - F_2$$

$$= \frac{\mu_0 I_1 I_2}{2\pi} - \frac{\mu_0 I_1 I_2}{4\pi}$$

$$= \frac{\mu_0 I_1 I_2}{4\pi} \quad \text{to right Repulsive}$$

Q.15 A force of 10N acts on a body of mass 20kg for 10 sec

Option 1:

50 kg m / s

Option 2:

100 kg m / s

Option 3:

300 kg m / s

Option 4:

200 kg m / s

Correct Answer:

100 kg m / s

Solution:

As we learnt in

Linear Momentum -

Quantity of motion in a body

$$\vec{P} = M \times \vec{V}$$

M= Mass and V= velocity



\vec{P} = Linear Momentum

- wherein

* Vector Quantity

* Same as that of velocity


Unit- Kg m/sec (S.I)




$$\Delta \vec{P} = f \cdot \Delta t \Rightarrow \Delta P = 10 \times 10 = 100 \text{ kgms}^{-1}$$

Q.16 Question is based on the following paragraph.

Wave property of electrons implies that they will show diffraction. Germer demonstrated this by diffracting electrons from a crystal. Diffraction from a crystal is obtained by requiring that the waves from different planes of atoms in a crystal interfere constructively (see diagram).




Question : In an experiment, electrons are made to pass through a narrow slit of width comparable to their de Broglie wavelength. They are detected on a screen at a distance D from the slit.




Which of the following graphs can be expected to represent the intensity of electrons detected as a function of the detector position ($y = 0$ corresponds to the central maximum)?


Option 1:




Option 2:




Option 3:



Option 4:



Correct Answer:



Solution:

As we learnt in

Davisson Germer Experiment -

A beam of electrons emitted by the electron gun is made to fall at a particular angle.

The scattered electron is received by the detector which can be moved along the arc.

The electron beams at the edges of the slit will diffract away from the central line. The electron beam along or near the central line will not diffract. The diffraction pattern is wider than the slit width.

As number of electron decreases both side of central maximum.

Correct graph is 1.

Q.17 A solid sphere of radius r made of a soft material of bulk modulus K is placed in a cylindrical container. A massless piston of area a is placed on the liquid, covering entire cross section of cylindrical container. To compress the liquid, the fraction of decrease in radius of the sphere,

$$\left(\frac{dr}{r}\right), \text{ is:}$$

Option 1:

$$\frac{mg}{Ka}$$

Option 2:

$$\frac{Ka}{mg}$$

Option 3:

$$\frac{Ka}{3mg}$$

Option 4:

$$\frac{mg}{3Ka}$$

Correct Answer:

$$\frac{mg}{3Ka}$$

Solution:

As we learnt that

Bulk Modulus -

Ratio of normal stress to volumetric strain.

$$K = \frac{f/A}{-\Delta v/v} = \frac{-Fv}{A\Delta v}$$

$$K = \frac{-Pv}{\Delta v}$$


v = Original volume

ΔV = Change in volume

P = Increase in pressure

-ve (sign) should be taken as decrease.

- wherein



$$\Delta P = \frac{mg}{a} \quad V = \frac{4\pi}{3}r^3$$

$$K = -\frac{\Delta P}{\left(\frac{\Delta V}{V}\right)} \quad \therefore \frac{dV}{V} = 3 \cdot \frac{dr}{r}$$

$$K = \frac{-\Delta P}{3 \cdot \left(\frac{dr}{r}\right)}$$

$$\text{or } \frac{dr}{r} = \frac{-mg}{3Ka}$$

Q.18 A uniform disc of mass 2 kg is rotated about an axis perpendicular to its plane and passing through its center. If the radius of gyration is 50 cm, then the M.I. of disc about this axis is

Option 1:

$$0.25 \text{ kg}^2$$

Option 2:

$$0.5 \text{ kg}^2$$

Option 3:

$$2 \text{ kg}^2$$

Option 4:

$$1 \text{ kg}^2$$

Correct Answer:

$$0.5 \text{ kg}^2$$

Solution:

As we learnt in

Radius of gyration (K) -

Radius of Gyration of a body about an axis is the effective distance from the axis at which the mass can be assumed to be concentrated so that moment of inertia remains the same.

- wherein

$$I = MK^2$$

$$K = \sqrt{\frac{I}{M}}$$

$$\text{use } I = MK^2$$

$$= 2 \times 50 \times 50 = 5000 \text{ kgcm}^2$$

$$= 0.5 \text{ kgm}^2$$

Q.19 What is the mechanical advantage of single fixed pulley?

Option 1:

1

Option 2:

2

Option 3:

0.5

Option 4:

4


Correct Answer:

1

Solution:

As we learnt in

Motion of connected blocks over pulley -



Equation of motion for

$$F_{net} = T - m_1g = m_1a$$

Equation of Motion for m_2

$$F_{net} = m_2g - T = m_2a$$

- wherein

$$a = \frac{[m_2 - m_1]g}{m_1 + m_2}$$

$$T = \frac{2m_1m_2 g}{m_1 + m_2}$$

Q. 20 A current of 2 mA was passed through an unknown resistor. Dissipated power when an ideal power supply of 11 V

Option 1:

$$11 \times 10^{-3} \text{ W}$$

Option 2:

$$11 \times 10^{-5} \text{ W}$$

Option 3:

$$11 \times 10^5 \text{ W}$$

Option 4:

$$11 \times 10^{-4} \text{ W}$$

Correct Answer:

$$11 \times 10^{-5} \text{ W}$$

Solution:

$$I = 2 \times 10^{-3} \text{ A}, P = 4.4 \text{ W}$$

$$P_{initial} = P = I^2 R \Rightarrow R = \frac{P}{I^2}$$

$$R = \frac{4.4}{4 \times 10^{-6}} = 1.1 \times 10^6 \Omega$$

$$P_{final} = P' = \frac{V^2}{R} = \frac{11^2}{R} = 11 \times 10^{-5} \text{ W}$$

Hence, (2) is the correct option.

Q.21 Dispersive power of crown glass is:

Option 1:

0.1639

Option 2:

0.1821

Option 3:

0.1764

Option 4:

0.2123

Correct Answer:


0.1639

Solution:

As we learn

Dispersive power (

$$\omega = \frac{\mu_v - \mu_r}{\mu_y - 1}$$



- wherein

μ_v = Refractive index of violet

μ_r = Refractive index of red

μ_y = Refractive index of yellow


$$= \frac{\mu_v + \mu_r}{2}$$

$$w = \frac{\mu_v - \mu_r}{\mu_y - 1}$$

$$\mu_y = \frac{\mu_v + \mu_r}{2} = 1.5187$$

$$\therefore w = \frac{1.523 - 1.5145}{1.5187 - 1} = 0.1639$$

Q. 22 A small ball of mass m starts at a point A . with speed v_0 on a smooth track AB as shown. The track BC has coefficient of friction μ after travelling a distance L which is :



Option 1:

$$\frac{2h}{\mu} + \frac{v_0^2}{2\mu g}$$

Option 2:

$$\frac{h}{\mu} + \frac{v_0^2}{2\mu g}$$

Option 3:

$$\frac{h}{2\mu} + \frac{v_0^2}{\mu g}$$

Option 4:

$$\frac{h}{2\mu} + \frac{v_0^2}{2\mu g}$$

Correct Answer:

$$\frac{h}{\mu} + \frac{v_0^2}{2\mu g}$$

Solution:

Solution :

Given :

Initial speed = v_0

Initial height = h

Applying conservation of energy,

$$mgh + \frac{1}{2}mv_0^2 = \frac{1}{2}mv^2$$

$$v^2 = v_0^2 + 2gh$$

Now, if a be the deceleration produced because of friction

Using,

Newton's third equation of motion,

$$v^2 = u^2 + 2as$$

$$0 = v_0^2 + 2gh - 2aL$$

$$L = \frac{v_0^2 + 2gh}{2a} \quad - (1)$$

Now, $f = \mu N$

where, $N = mg$

$$f = \mu mg$$

$$a = \mu g$$

Putting back this value in equation (1)

$$\begin{aligned} L &= \frac{v_0^2 + 2gh}{2\mu g} \\ &= \frac{h}{\mu} + \frac{v_0^2}{2\mu g} \end{aligned}$$

-
- Q. 23 A moving coil galvanometer, having a resistance G , produces a current I_g flows through it. This galvanometer can be converted into a voltmeter ($I_0 > I_g$) by connecting a series resistance R_A and into a ammeter ($I_0 < I_g$) by connecting a shunt resistance R_V in parallel to it.

Option 1:

$$R_A R_V = G^2 \left(\frac{I_0 - I_g}{I_g} \right)^2 \text{ and } \frac{R_A}{R_V} = \left(\frac{I_g}{(I_0 - I_g)} \right)^2$$

Option 2:

$$R_A R_V = G^2 \text{ and } \frac{R_A}{R_V} = \left(\frac{I_g}{I_0 - I_g} \right)^2$$

Option 3:

$$R_A R_V = G^2 \left(\frac{I_g}{I_0 - I_g} \right) \text{ and } \frac{R_A}{R_V} = \left(\frac{I_0 - I_g}{I_g} \right)^2$$

Option 4:

$$\frac{R_A}{R_V} = G^2 \text{ and } \frac{R_A}{R_V} = \frac{I_g}{(I_0 - I_g)}$$

Correct Answer:


$$R_A R_V = G^2 \text{ and } \frac{R_A}{R_V} = \left(\frac{I_g}{I_0 - I_g} \right)^2$$

Solution:

Conversion of galvanometer into voltmeter -

Connected a large resistance

- wherein



$$(I_0 - I_G)R_A = I_G G \quad V_0 = I_G(G + R_A)$$

$$V_0 = I_0 G$$

$$R_A = \frac{I_G G}{I_0 - I_G}$$

$$I_0 G = I_G(G + R_V)$$


$$\frac{I_0 G}{I_G} - G = R_V$$

$$R_V = \frac{G(I_0 - I_G)}{I_G}$$

$$\frac{R_A}{R_V} = \frac{I_G G}{I_0 - I_G} \times \frac{I_G}{G(I_0 - I_G)} = \left(\frac{I_G}{I_0 - I_G} \right)^2$$

$$R_A R_V = \frac{I_G G}{I_0 - I_G} \times \frac{G(I_0 - I_G)}{I_G} = G^2$$

Q. 24 The V-I characteristic of a diode is shown in the figure. resistance is :



Option 1:

10

Option 2:

10

Option 3:

106

Option 4:

100

Correct Answer:

10

Solution:

Forward Resistance

$$R_F = \frac{\Delta V}{\Delta i} = \frac{0.1}{10 \times 10^{-3}} \Omega$$

$$\text{Reverse bias Resistance} = \frac{\Delta V}{\Delta i} = \frac{10}{10^{-6}} = 10^7 \Omega$$

Ratio of forward to reverse resistance = $\frac{10}{10^7} = 10^{-6}$

Correct option is 2.

Q.25 The current voltage relation of diode is relation of diode where the applied voltage V is in volts and the temperature makes an error $\pm 0.01\%$ while measuring the current of 5 mA. be the error in the value of current in mA ?

Option 1:

0.2 mA

Option 2:

0.02 mA

Option 3:

0.5 mA

Option 4:

0.05 mA

Correct Answer:

0.2 mA

Solution:

Given $I = 5 \text{ mA} = \left(e^{\frac{1000V}{T}} - 1 \right)$

or

$$e^{\frac{1000V}{T}} = 6$$

Also $I = e^{\frac{1000V}{T}} - 1$

$$\frac{dI}{dV} = \left(\frac{1000}{T} \right) \cdot e^{\frac{1000V}{T}}$$

$$\Rightarrow dI = \left(e^{\frac{1000V}{T}} \right) \cdot \frac{1000}{T} dV$$

$$\Rightarrow dI = 6 \times \frac{1000}{300} \times 10^{-2}$$

$$= 0.2 \text{ mA}$$

Q.26 Work done by static friction on an object:

Option 1:

may be positive

Option 2:

must be negative

Option 3:

must be zero

Option 4:

none of these

Correct Answer:

may be positive

Solution:

Work done by the frictional force is negative -


When the force is large enough to overcome the friction

Work done by the frictional force is positive -

When force is applied on a body, which is placed above another body, the frictional force on the lower body may be positive


Work done by the frictional force is zero -

When force applied on a body is insufficient to overcome the friction



Consider the blocks shown in the figure to be moving together due to friction between them.

The free body diagrams of both the blocks are shown below.



Work done by static friction on A is positive and on B is negative.

Q. 27 A green light is incident from the water to the air - water interface. Select the correct statement.

Option 1:

The entire spectrum of visible light will come out of the water.

Option 2:

The spectrum of visible light whose frequency is less than that of the denser medium.

Option 3:

The spectrum of visible light whose frequency is more than that of the denser medium.

Option 4:

The entire spectrum of visible light will come out of the water.

Correct Answer:

The spectrum of visible light whose frequency is less than that of the denser medium.

Solution:


As we learnt in

Critical angle -

$\sin c = \frac{\text{Refractive index of rarer medium}}{\text{Refractive index of denser medium}}$

- wherein

When angle of incidence of a ray is equal to the critical angle, no refraction occurs.



Also refractive index (μ) of the medium depends on the wavelength. So will be move for lenser frequency of light.

Correct option is 3.

Q. 28 Match List I (Wavelength range of 22. electromagnetic production of these waves) and select the correct options.

List I		List II	
(a)	700 nm to 1 mm	(i)	Vibration of atoms and molecules.
(b)	1 nm to 400 nm	(ii)	Inner shell electrons in atoms moving from one energy level to a lower level.
(c)	$< 10^{-3}$ nm	(iii)	Radioactive decay of the nucleus.
(d)	1 mm to 0.1 m	(iv)	Magnetron valve.

Option 1:

(a) - (iv), (b) - (iii), (c) - (ii), (d) - (i)

Option 2:

(a) - (iii), (b) - (iv), (c) - (i), (d) - (ii)

Option 3:

(a) - (ii), (b) - (iii), (c) - (iv), (d) - (i)

Option 4:

(a) - (i), (b) - (ii), (c) - (iii), (d) - (iv)

Correct Answer:


(a) - (i), (b) - (ii), (c) - (iii), (d) - (iv)

Solution:

Magnetic flux -

The total number of magnetic lines of force passing normally

- wherein



700 nm to 1 mm \rightarrow Vibration of atoms and molecule

1 nm to 400 nm \rightarrow inner shell electrons moving from one energy level

$< 10^{-3}$ nm (r radius of α particle) \rightarrow Radioactive decay of nucleus

1 mm to \rightarrow 1 m \rightarrow Magnetron valve.

Q.29 One mole of an ideal gas when went through a thermody
volume changed from p and V to 2P and 2V then its entr

Option 1:

$$C_v \log_e 2$$

Option 2:

$$C_p \log_e 2$$

Option 3:

$$(C_p + C_v) \log_e 2$$

Option 4:

$$(C_p - C_v) \log_e 2$$

Correct Answer:

$$(C_p + C_v) \log_e 2$$

Solution:

As we have learned

Entropy change in terms of Pressure and Volume -

$$\Delta S = n c_v \ln \left(\frac{P_2}{P_1} \right) + n C_P \ln \left(\frac{V_2}{V_1} \right)$$

-

entropy change in term of P and V is

$$\Delta S = n c_v \ln \left(\frac{P_2}{P_1} \right) + n C_P \ln \left(\frac{V_2}{V_1} \right)$$

$$\Delta S = n c_v \ln \left(\frac{2P}{P_1} \right) + n C_P \ln \left(\frac{2V}{V_1} \right)$$

$$\Delta S = c_v \log_e(2) + C_P \log_e(2)$$

$$\Delta S = \log_e(2) \times (C_v + C_P)$$

- Q.30 This question has statement 1 and statement 2 Of the following statements, choose the one that best describes the two statements.
- statement 1 : Davisson – germer experiment established wave nature of electrons
- statement 2 : If electrons have wave nature, they can interfere

Option 1:

Statement 1 is false, Statement 2 is true

Option 2:

Statement 1 is true, Statement 2 is false

Option 3:

Statement 1 is true, Statement 2 is true and the correct explanation is given for statement 2.

Option 4:

Statement 1 is true, Statement 2 is true, Statement 2 is not explained for statement 1.

Correct Answer:

Statement 1 is true, Statement 2 is true and the correct explanation is given for statement 2.

Solution:

Davisson-Germer experiment showed that electron beams can diffract from atomic crystals. This shows the wave nature of electrons as wave diffraction.


Correct option is

Any wave will show interference and diffraction and interference during Davisson-Germer experiment hence it establishes wave nature of electrons.


Chemistry

Q.1 Which one of the following would not be oxidised upon treatment with $\text{K}_2\text{Cr}_2\text{O}_7$ and H_2SO_4 ?


Option 1:




Option 2:




Option 3:



Option 4:




Correct Answer:



Solution:

As we learnt,

Tertiary alcohols do not get oxidised in presence of H_2SO_4 and $\text{K}_2\text{Cr}_2\text{O}_7$ it undergoes elimination reaction and alkenes are produced.



Therefore option (3) is correct.

Q.2 The Coagulation of colloidal particles of the sol can be

Option 1:

Heating

Option 2:

Adding oppositely charged sol

Option 3:

Adding Electrolyte

Option 4:

All of above

Correct Answer:

All of above

Solution:

By mixing two oppositely charged sols: Oppositely charged sol neutralise their charges and get partially or completely precipitated (e.g. ferric hydroxide (+ve sol) and arsenious sulphide (-ve sol) bring them in the precipitate called mutual coagulation.

By boiling/heating: When a sol is boiled, the adsorbed layer is destroyed, and the molecules of dispersion medium. This reduces the charge on the particles, causing them to settle down in the form of a precipitate.

By addition of electrolytes: When excess of an electrolyte is added to a sol, it causes coagulation. The reason is that colloids interact with ions carrying charge. The ion responsible for coagulation is called the coagulating ion. A negative ion causes coagulation of a positive sol and vice versa.

So, all are correct.

Therefore, option number (4) is correct.

Q.3 The tungsten filament for electric bulb is formed by

Option 1:

Graphite

Option 2:

Diamond

Option 3:

F u l l e r e n e

Option 4:

C h a r c o a l

Correct Answer:

D i a m o n d

Solution:

A s w e h a v e l e a r n t ,

D i a m o n d i s u s e d f o r s h a r p e n i n g h a r d t o o l s , i n m a k i n g d i e s , j e w e l l e r y b u l b s .

Therefore, option (2) is correct.

Q. 4 Which of the following has highest metallic character?

Option 1:

P

Option 2:

A s

Option 3:

S b

Option 4:

B i

Correct Answer:

B i

Solution:

O n m o v i n g d o w n t h e g r o u p , t h e m e t a l l i c c h a r a c t e r i n c r e a s e s .

N i t r o g e n a n d p h o s p h o r u s a r e n o n - m e t a l s , A r s e n i c a n d a n t i m o n y a r e m e t a l l o i d s .

Therefore, **option (4) is correct.**

Q.5

A n e l e m e n t h a s t h e e l e c t r o n c o n f i g u r a t i o n $1s^2, 2s^2, 2p^6, 3s^2, 3p^2$. To which block in the periodic table, does this belong?

Option 1:

s - b l o c k

Option 2:

p - b l o c k

Option 3:

d - b l o c k

Option 4:

f - b l o c k

Correct Answer:

p - b l o c k


Solution:

A s w e l e a r n t i n

A u f b a u P r i n c i p l e -

In the ground state of the atoms, the orbitals are filled in order of their increasing energies.

- w h e r e i n



According to Aufbau's principle, in ground state of atoms, the orbitals are filled in order of their increasing energies. Clearly, with unfilled 3p orbitals, the elements in this block are the p-block elements.

Q.6 Which of the following has highest boiling point?

Option 1:

ICl

Option 2:

BrCl

Option 3:

Br_2

Option 4:

Cl_2

Correct Answer:

ICl

Solution:

As we have learnt,

Interhalogen compounds have higher boiling point than Halogens.

Boiling Points depends on bond polarity and molecular mass. The greatest molecular mass along with the greatest bond polarity.

The correct order of boiling point is given as

$\text{ICl} > \text{BrCl} > \text{Br}_2 > \text{Cl}_2$

Hence the correct answer is Option (1)

Q. 7 Which of the following do not impart any colour to flame?

Option 1:

Be

Option 2:

Al

Option 3:

Both a and b

Option 4:

Ca

Correct Answer:

Both a and b

Solution:

As we learn

Colour of flame due to Be and Al -


Both do not impart colour to flame

-

Be and Al both do not impart colour to flame.

Q.8 Perhydrol is the name associated with

Option 1:



Option 2:




Option 3:



Option 4:



Correct Answer:




Solution:

As we learnt in

Hydrogen Peroxide -

Discovered by J - L Thenard

- wherein



Perhydrol is a 30% weight by volume solution of hydrogen peroxide

Q.9 Which of the following hydride has the lowest boiling point

Option 1:

N H 3

Option 2:

P H 3

Option 3:

S b H 3

Option 4:

A s H 3

Correct Answer:

P H 3

Solution:

From PH₃ to BiH₃ the boiling points increase with the increase in Van der Waal's forces with molecular mass. However, the b has a lower molecular mass than PH₃, due to the intermolecular forces not present in PH₃ and other hydrides in Group-15.

Therefore, Option(2) is correct

Q.10

The bond order in NO is 2.5 while that in NO⁺ is 3. Which has greater bond length for these two species?

Option 1:

Bond length is greater than in NO⁺

Option 2:

Bond length is greater than in NO⁺

Option 3:

Bond length is unequal to that in NO⁺

Option 4:

Bond length is unpredictable.

Correct Answer:

Bond length is greater than in NO⁺

Solution:

Bond order is defined as one half the difference between the number of bonding and the antibonding orbitals.


$$\text{Bond Order} = \frac{N_b - N_a}{2}$$

The higher the bond order, the shorter will be the bond length. NO^+ has a shorter bond length than that of NO so NO^+ has a shorter bond length.


Q.11 The organic compound that gives following qualitative :

Test	Inference
1. HCl	Insoluble
2. NaOH solution	soluble
3. Br_2/water	Decolourization


Option 1:




Option 2:




Option 3:



Option 4:




Correct Answer:



Solution:

Physical properties of Phenol -

- Phenols are more acidic than alcohols
- They are less soluble in water but readily soluble in organic solvents
- They liquify due to high hygroscopic nature.



The re ~~foptione~~ (1) is correct.

Q. 12 Select the incorrect option regarding metallic solids

Option 1:

Positive ions are surrounded by sea of free electrons

Option 2:

These electrons are mobile

Option 3:

Covalent bonds are strong and directional

Option 4:

These are highly malleable and ductile

Correct Answer:

Covalent bonds are strong and directional

Solution:

As we learnt in

Metallic solids -

Metallic bond, i.e., attraction between positively charged me

- wherein

Ex. iron, copper, zinc, aluminium, sodium

In metallic solids positive ions are surrounded by free elecvt
are highly ductile and malleable. Covalent bonds are strong a
not in metallic solid.

Q.13 Pt and Pd adsorb large volumes of hydrogen under spec adsorbed is called?

Option 1:

Adsorbed Hydrogen

Option 2:

Reductive hydrogen

Option 3:

Occluded hydrogen

Option 4:

Atomic hydrogen

Correct Answer:

Occluded hydrogen

Solution:

Adsorption of any gas on the metal surface is also sometimes adsorbed on a metal surface, then this hydrogen is known as

Hence, the correct answer is Option (3)

Q.14

Consider the following table :

Gas	$a/(k Pa dm^6 mol^{-1})$	$b/(dm^3 mol^{-1})$
A		642.32
B		155.21
C		431.91
D		155.21

a and b are van der Waals constants. The gas

Option 1:

Gas C will occupy more volume than gas A;
gas B will be more compressible than gas D

Option 2:

Gas C will occupy lesser volume than gas A;
gas B will be lesser compressible than gas D

Option 3:

Gas C will occupy more volume than gas A;
gas B will be lesser compressible than gas D

Option 4:

Gas C will occupy
lesser volume than gas
A; gas B will be more
compressible than gas

Correct Answer

Gas C will occupy more volume than gas A;
gas B will be more compressible than gas D

Solution:

Vander Waal equation for real gas -

$$\left(p + \frac{an^2}{v}\right)(V - nb) = nRT$$

Vander Waal constant 'a' is the measure of the intermolecular attraction of a gas.

Vander Waal constant 'b' is a measure of the size of the molecule.

Greater the value of 'a', greater is the force of attraction &

Greater the value of 'b', the higher will be the effective vol

Gas A and C have same value of 'b' but different value of 'a' so
force of attraction so molecules will be more closer hence occ

Gas B and D have same value of 'a' but different value of 'b' so
compressible.

∴ Option (1) is correct.

Q.15 The electrons identified by quantum numbers n and l :

(1) n = 4, l = 1 (2) n = 4, l = 0 (3) n = 3, l = 2 (4) n = 3, l = 1

Can be placed in order of increasing energy as :

Option 1:

(4) < (2) < (3) < (1)

Option 2:

$$(2) < (4) < (1) < (3)$$

Option 3:

$$(1) < (3) < (2) < (4)$$

Option 4:

$$(3) < (4) < (2) < (1)$$

Correct Answer:

$$(4) < (2) < (3) < (1)$$

Solution:

As discussed in the concept
Principal Quantum Number (n) -

It is a positive integer with value of $n = 1, 2, 3, \dots$

-

and

Azimuthal Quantum Number (l) -

For a given value of n, l can have n values ranging from 0 to n - 1, that is, for a given value of n, the possible value of l are : $l = 0, 1, 2, \dots, (n-1)$

-

$$(1) \ n = 4, l = 1 \Rightarrow 4p$$

$$(2) \ n = 4, l = 0 \Rightarrow 4s$$

$$(3) \ n = 3, l = 2 \Rightarrow 3d$$

$$(4) \ n = 3, l = 1 \Rightarrow 3p$$

Increasing order of energy is

$$3p < 4s < 3d < 4p$$

$$(4) < (2) < (3) < (1)$$

Alternatively,

$$\text{For } (1) \ n + l = 5; n = 4$$

$$(2) \ n + l = 4; n = 4$$

$$(3) \ n + l = 5; n = 3$$

$$(4) \ n + l = 4; n = 3$$

Lower $n + l$ means less energy and if for two subshells $n + l$ is same than lower n , lower will be the energy.

Thus correct order is $(4) < (2) < (3) < (1)$

Q.16 Which of following is not an ambidentate ligand?

Option 1:



Option 2:



Option 3:



Option 4:



Correct Answer:




Solution:


As we learnt in

Types of Ligands on the basis of Connectivity -

(i) Unidentate - ligand is bound to metal ion through a single



(ii) Bidentate : when ligand can bind through two donor atoms



(iii) Polydentate - when ligand bind to two or more donor atoms

(iv) Hexadentate - type of polydentate having six donor atoms

(v) Ambidentate - which can ligate through two different atoms



- wherein

All ambidentate ligands are monodentate but all monodentate

Ambidentate Ligands can ligate through two different atoms and
Of all the given, CO is the only ligand which can donate only

Q.17 The edge length of a face centred cubic cell of an ionic
the cation is 110 pm, the radius of the anion is :

Option 1:

144 pm

Option 2:

288 pm

Option 3:

398 pm

Option 4:

618 pm

Correct Answer:

144 pm

Solution:

As we learnt in

Centered unit cell -

In centered unit cell, constituent particles are present in positions
corners.

- wherein


Three types: face centered, body centered, end centered.

For fcc unit cell, $2(\text{radius of cation} + \text{radius of anion}) = \text{edge length}$


$$\text{Therefore } (2r_c + r_a) = \frac{508}{2} \Rightarrow r_a = 254 - 110 = 144 \text{ pm}$$

Q.18 Chlorine on reaction with hot and concentrated sodium


Option 1:




Option 2:




Option 3:



Option 4:




Correct Answer:



Solution:


Reaction of chlorine with hot and concentrated NaOH

Cl_2 gives disproportionation reaction on reaction with hot and concentrated NaOH to give Chloride and Chlorate ions




Therefore **Option (4)** is correct


Q.19 The major product expected from the following reaction




Option 1:




Option 2:



Option 3:



Option 4:



Q. 20 XeF_6 cannot be stored in glass because

Option 1:

it doesn't react

Option 2:

It reacts with SiO_2

Option 3:

It reacts with HCl

Option 4:

Both a and c

Correct Answer:

It reacts with SiO_2


Solution:

As we learnt in




Structure of Xenon hexa uoride -

Distorted octahedral and hybridisation is Sp^3d^3

- wherein



XeF_6 cannot be stored in glass because it reacts with the silicic acid



Q. 21 The standard reduction potential of four element are A = 0.68 V, B = 0.15 V, C = 0.77 V, D = -0.4202 V

The element that displaces A from its compounds is

Option 1:

B

Option 2:

C

Option 3:

D

Option 4:

none of these

Correct Answer:

D

Solution:

As we have learned

Standard Electrode Potential -

If concentration of each species taking part in the electrode reaction is 1 M and the reaction is carried out at 298 K, then the potential of each electrode is standard electrode potential.

A is displaced from D because D was an

Q.22 18 g of glucose is added to 178.2 g of water. The vapour pressure of this aqueous solution at 100°C is

Option 1:

759.00 torr

Option 2:

7.60 torr

Option 3:

76.00 torr

Option 4:

752.40 torr.

Correct Answer:

752.40 torr.

Solution:

As we learnt in

Expression of relative lowering of vapour pressure -

$$\frac{\Delta P}{P^0} = x_{\text{solute}}$$

$$x_{\text{solute}} = \frac{n_{\text{solute}}}{n_{\text{solute}} + n_{\text{solvent}}}$$

ΔP is lowering of v.p.

$P^0 \rightarrow$ vapour pressure of pure solvent

$x_{\text{solute}} \rightarrow$ mole fraction of non volatile solute

$$\frac{P_A^0 - P_s}{P_A^0} = \chi_B$$

$$\text{Mole fraction of glucose} = \frac{18/180}{\frac{18}{180} + \frac{178.2}{18}} = \frac{1}{100}$$

$$P_A^0 = \text{Vapour pressure of pure water at } 100^\circ\text{C} = 760 \text{ torr}$$

$$\frac{760 - p_s}{760} = \frac{1}{100}$$

p_s = Vapour pressure of solution = 752.4 torr.

Correct option is 4.

Q.23 Oxidation Number of S in H_2SO_5

Option 1:

+ 6

Option 2:

+ 8

Option 3:

+ 4

Option 4:

+ 2


Correct Answer:

+ 6

Solution:

As we have learnt,

The structure of Peroxomonosulphuric acid is given below:



From the structure we can say S has +6 oxidation state.


Therefore option (1) is correct.

Q.24 Which of the following is (are) primary standard?


Option 1:

$\text{Na}_2\text{C}_2\text{O}_4$

Option 2:



Option 3:



Option 4:

All of these

Correct Answer:

All of these

Solution:

As we learnt

Primary Standard -

Primary standard is a substance of known high purity which n solvent to give a primary standard solution. It is a reference concentration of another known chemical.

- wherein

NaC_2O_4 (Sodium Oxalate) ; $K_2Cr_2O_7$ (Potassium Dichromate) ; KIO_3 (Potassium Iodate) ; Oxalic acid

All of the given chemicals are primary standard

Q.25 Which of the following have least density -

Option 1:

C

Option 2:

Si

Option 3:

Ge

Option 4:

Sn

Correct Answer:

Si

Solution:

Fact based.

Order of density of the elements is $\text{Sn} > \text{Ge} > \text{C} > \text{Si}$

Therefore, option (2) is correct.

Q. 26 Helium atom is two times heavier than a Hydrogen molecule. The average translational energy of Helium is

Option 1:

Two times that of a hydrogen molecule

Option 2:

Same as that of a hydrogen molecule

Option 3:

Four times that of a hydrogen molecule

Option 4:

Half that of a hydrogen molecule

Correct Answer:

Same as that of a hydrogen molecule

Solution:

Average translational energy per molecule of any ideal gas is


$$E = \frac{3kT}{2}$$

$$\frac{E_{\text{He}}}{E_{\text{H}_2}} = \frac{T_{\text{He}}}{T_{\text{H}_2}}$$

So energies will be same for He and H₂ at same temperature.

Hence, the correct answer is Option (2)

Q.27 Increasing S_N1 rate of reaction in the following compounds is :



Option 1:

(A) < (B) < (C) < (D)

Option 2:

(B) < (A) < (C) < (D)

Option 3:

(B) < (A) < (D) < (C)

Option 4:

(A) < (B) < (D) < (C)


Correct Answer:

(B) < (A) < (C) < (D)

Solution:

As we have learnt,

Rate of S_N1 is directly proportional to stability of first formed car



∴ increasing S_N1 rate reaction : (B) < (A) < (C) < (D)

Therefore, option (2) is correct.

Q. 28 Shape of s orbital is:

Option 1:

Spherical

Option 2:

Dumb bell

Option 3:

Cylindrical

Option 4:

Elliptical

Correct Answer:

Spherical

Solution:

As we learnt

Shape of s orbital -

spherical

-

s– orbital ($l=0$) have no directional dependence and are spherically symmetrical.

Therefore, **Option(1) is correct**

Q. 29 Which one of the following aqueous solutions will exhibit

Option 1:

$0.01M Na_2SO_4$

Option 2:

$0.01M KNO_3$

Option 3:

$0.015M Urea$

Option 4:

$0.015M glucose$

Correct Answer:

$0.01M Na_2SO_4$

Solution:

Elevation in Boiling point $\propto i \times m$

For Na_2SO_4 , $i = 3$, will be highest among these given aqueous

So, In $0.01M Na_2SO_4$ it will be 0.03 which is the highest in the

The correct option is 1.

Q. 30 Which of the following ligand is bidentate?

Option 1:

$C_2O_4^{2-}$


Option 2:

$CH_3C \equiv N$


Option 3:

Br^-

Option 4:



Correct Answer:




Solution:


As we learnt in

Types of Ligands on the basis of Connectivity -

(i) Unidentate - ligand is bound to metal ion through a single




(ii) Bidentate : when ligand can bind through two donor atoms



(iii) Polydentate - when ligand binds to two or more donor atoms

(iv) Hexadentate - type of polydentate having six donor atoms

(v) Ambidentate - which can ligate through two different atoms



- wherein

All ambidentate ligands are monodentate but all monodentate

When Ligands can bind through 2 donor atoms they are called $C_2O_4^{2-}$ is that criteria.

Maths

Q.1

The sum $\sum_{k=1}^{20} (1 + 2 + 3 + \dots + k)$ is

Option 1:

1540

Option 2:

1680

Option 3:

1 2 6 0

Option 4:

1 4 5 0

Correct Answer:

1 5 4 0

Solution:

Sum of Common Series

Sum of the square of first n-natural numbers

$$1^2 + 2^2 + 3^2 + 4^2 + + n^2 = \frac{n(n+1)(2n+1)}{6}$$

Now ,

$$\begin{aligned}\sum_{k=1}^{20} \frac{k(k+1)}{2} &= \frac{1}{2} \left(\sum_{k=1}^{20} k^2 + \sum_{k=1}^{20} k \right) \\ &= \frac{1}{2} \left(\frac{20(20+1)(2 \times 20 + 1)}{6} + \frac{20(20+1)}{2} \right) \\ &= \frac{1}{2} ((70 \times 41) + 210) = 1540\end{aligned}$$

Q.2 What is the symbol of OR disjunction?

Option 1:

U

Option 2:

∩

Option 3:

∪

Option 4:

Both (a) and (c)

Correct Answer:

Both (a) and (c)

Solution:

As we have learned

Disjunction "OR" -

The symbol for "OR" disjunction is the symbol

Q.3 The plane $2x - z = 4$ cuts the sphere $x^2 + y^2 + z^2 - x + z - 2 = 0$ in a circle of radius

Option 1:

1

Option 2:

3

Option 3:

$\sqrt{2}$

Option 4:

2

Correct Answer:

1

Solution:

Equation of Sphere -

Equation of the sphere is

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$$

NOTE:

If the centre of the sphere at origin the equation of sphere is

The equation of the sphere can also be written as $x^2 + y^2 + z^2 + 2(ax + by + cz) + d = 0$, where $d = a^2 + b^2 + c^2 - r^2$

-

Centre $\left(\frac{1}{2}, 0, -\frac{1}{2}\right)$ $r = \sqrt{\frac{5}{2}}$

Perpendicular distance from centre = $\frac{\left|\frac{1}{2} + \frac{1}{2} - 4\right|}{\sqrt{6}} = \sqrt{\frac{3}{2}}$

$$\text{Radius}^2 = \left(\sqrt{\frac{5}{2}}\right)^2 - \left(\sqrt{\frac{3}{2}}\right)^2 = 1$$

Q.4 Equation of curve such that length of $\sqrt{2}$ times to dd inwards to that point, will be

Option 1:

$$y = 2x + c$$

Option 2:

$$y = x + c$$

Option 3:

$$y = 3x + c$$

Option 4:

$$y = 4x + c$$

Correct Answer:

$$y = x + c$$


Solution:

AS we have learnt,

Length of Tangent -

$$\text{Length} = \left| y \sqrt{1 + \left(\frac{dx}{dy} \right)^2} \right|$$

- wherein



$$\left| y \sqrt{1 + \left(\frac{dx}{dy} \right)^2} \right| = \sqrt{2}y$$

On squaring, we get

$$y^2 \left(1 + \left(\frac{dx}{dy} \right)^2 \right) = 2y^2 \Rightarrow \frac{dx}{dy} = 1, -1$$

$$\Rightarrow y = x + c \text{ or } y = -x + c$$

Q. 5 In ΔABC $\cot \frac{A}{2} \cdot \cot \frac{B}{2} \cdot \cot \frac{C}{2}$ is equal to

Option 1:

$$\frac{\Delta}{r^2}$$

Option 2:

$$\frac{(a+b+c)^2}{abc} \cdot 2R$$

Option 3:

$$\frac{\Delta}{r}$$

Option 4:

$$\frac{\Delta}{Rr}$$

Correct Answer:

$$\frac{\Delta}{r^2}$$

Solution:


Trigonometric Ratios of Functions -

$$\operatorname{cosec} \Theta = \frac{Hyp}{Opp}$$

$$\sec \Theta = \frac{Hyp}{Base}$$

$$\cot \Theta = \frac{Base}{Opp}$$

- wherein



$$\begin{aligned}
 & \cot \frac{A}{2} \cdot \cot \frac{B}{2} \cdot \cot \frac{C}{2} \\
 &= \frac{s(s-a)}{\Delta} \cdot \frac{s(s-b)}{\Delta} \cdot \frac{s(s-c)}{\Delta} \\
 &= \frac{s^3(a+b+c)}{\Delta^3} = \frac{s^2 \Delta^2}{\Delta^3} = \frac{s^2}{\Delta} \\
 &= \left(\frac{\Delta^2}{r^2} \right) \times \frac{1}{\Delta} = \frac{\Delta}{r^2} \quad \left[\because s = \frac{\Delta}{r} \right] \\
 & \cot \frac{A}{2} \cdot \cot \frac{B}{2} \cdot \cot \frac{C}{2} = \frac{\Delta}{r^2}
 \end{aligned}$$

Q. 6 If $A \cup B = P, A \cup C = Q, A \cap B = R, A \cap C = S, B \cup C = T, B \cap C = V$. Then simplify $A \cup (B \cap C)$

Option 1:

$$R \cap S$$

Option 2:

$$T \cap V$$

Option 3:

$$P \cap Q$$

Option 4:

$$P \cup Q$$

Correct Answer:

$$P \cap Q$$

Solution:

As we learnt

DISTRIBUTIVE PROPERTY -

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

-

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C) = P \cap Q$$

Q.7 The value of $\frac{1}{\cos 285^\circ} + \frac{1}{\sqrt{3} \sin 255^\circ}$, is

Option 1:

$$\sqrt{3} - \sqrt{2}$$

Option 2:

$$2\sqrt{2}$$

Option 3:

$$\frac{4\sqrt{2}}{\sqrt{3}}$$

Option 4:

$$\frac{2\sqrt{2}}{3}$$

Correct Answer:

$$\frac{4\sqrt{2}}{\sqrt{3}}$$

Solution:

$$\frac{1}{\cos 285^\circ} + \frac{1}{\sqrt{3} \sin 255^\circ}$$

$$= \frac{1}{\cos 75^\circ} + \frac{1}{\sqrt{3} (-\sin 75^\circ)} \quad \left[\begin{array}{l} \because \cos 285^\circ = +\cos 75^\circ \\ \sin 255^\circ = -\sin 75^\circ \end{array} \right]$$

$$= \frac{1}{\sin 15^\circ} - \frac{1}{\sqrt{3} \sin 75^\circ}$$

-

-

$$\begin{aligned}
 &= \frac{2\sqrt{2}}{\sqrt{3}-1} - \frac{1(2\sqrt{2})}{\sqrt{3}(\sqrt{3}+1)} \\
 &= 2\sqrt{2} \left[\frac{1}{\sqrt{3}-1} - \frac{1}{\sqrt{3}(\sqrt{3}+1)} \right] \\
 &= \frac{2\sqrt{2}}{\sqrt{3}} \frac{[(\sqrt{3}+1)\sqrt{3} - (\sqrt{3}-1)]}{2} = \frac{4\sqrt{2}}{\sqrt{3}}
 \end{aligned}$$

Q.8 The general solution of the differential equation

Option 1:

$$x^3y - 3e^x = cy$$

Option 2:

$$x^3y + 3e^x = 3cy$$

Option 3:

$$y^3x - 3e^y = cx$$

Option 4:

$$y^3x + 3e^y = cx$$

Correct Answer:

$$x^3y + 3e^x = 3cy$$

Solution:

As we learnt in

Linear Differential Equation -

$$\frac{dy}{dx} + Py = Q$$

- wherein

P, Q are functions of x alone.

$$y(x^2y + e^x)dx = e^x dy$$

$$\frac{1}{y^2} \frac{dy}{dx} - \frac{1}{y} = \frac{x^2}{e^x}$$

$$\text{Let } \frac{1}{y} = t$$

$$\therefore \frac{-1}{y^2} \frac{dy}{dx} = \frac{dt}{dx}$$

$$\therefore \frac{dt}{dx} + t = \frac{-x^2}{e^x}$$

$$\int 1 \cdot dx = x$$

$$I.F. = e^x$$

Solution is

$$t \cdot e^x = \int \frac{-x^2}{e^x} \times e^x dx$$

$$= \frac{-x^3}{3} + C$$

$$\frac{1}{y} e^x = \frac{-x^3}{3} + C$$

$$\therefore 3e^x = -x^3 y + 3cy$$

$$\therefore 3e^x + x^3 y = 3cy$$

Q. 9 $\lim_{x \rightarrow 3} \frac{\sqrt{3x} - 3}{\sqrt{2x - 4} - \sqrt{2}}$ is equal to

Option 1:

$$\sqrt{3}$$

Option 2:

$$\frac{1}{\sqrt{2}}$$

Option 3:

$$\frac{\sqrt{3}}{2}$$

Option 4:

$$\frac{1}{2\sqrt{2}}$$

Correct Answer:

$$\frac{1}{\sqrt{2}}$$

Solution:

Limit is of 0/0 form. We can rationalize the irrational powers

$$\begin{aligned}
& \lim_{x \rightarrow 3} \frac{\sqrt{3x} - 3}{\sqrt{2x - 4} - \sqrt{2}} \\
&= \lim_{x \rightarrow 3} \frac{\sqrt{3x} - 3}{\sqrt{2x - 4} - \sqrt{2}} \times \frac{\sqrt{2x - 4} + \sqrt{2}}{\sqrt{2x - 4} + \sqrt{2}} \times \frac{\sqrt{3x} + 3}{\sqrt{3x} + 3} \\
&= \lim_{x \rightarrow 3} \frac{3x - 9}{2x - 4 - 2} \times \frac{\sqrt{2x - 4} + \sqrt{2}}{\sqrt{3x} + 3} \\
&= \lim_{x \rightarrow 3} \frac{3(x - 3)}{2(x - 3)} \times \frac{\sqrt{2x - 4} + \sqrt{2}}{\sqrt{3x} + 3} \\
&= \frac{3}{2} \cdot \frac{\sqrt{2 \times 3 - 4} + \sqrt{2}}{\sqrt{3 \times 3} + 3} \\
&= \frac{3}{2} \times \frac{\sqrt{2} + \sqrt{2}}{6} \\
&= \frac{2\sqrt{2}}{4} = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}
\end{aligned}$$

Q.10 Find the Range

$$p = \frac{2q + 1}{q - 5}$$

Option 1:

$$p \neq 2$$

Option 2:

$$q \neq 5$$

Option 3:

$$p \neq 5$$

Option 4:

None of these.

Correct Answer:

$$p \neq 2$$

Solution:

As we learnt in

Range -

The range of the relation R is the set of all second elements

- wherein

e.g. $R = \{(a, b), (c, d)\}$. Then Range is $\{b, d\}$

$$p = \frac{2q + 1}{q - 5}$$

$$pq - 5p = 2q + 1$$

$$(p - 2)q = 5p + 1$$

$$q = \frac{5p + 1}{p - 2}$$

Here $p \neq 2$ is the range.

Q.11 Find the smallest positive root of the equation $\sqrt{\cos(1-x)} = \sqrt{\sin x}$

Option 1:

$$\frac{\pi}{4} - \frac{1}{2}$$

Option 2:

$$\frac{\pi}{4} + \frac{1}{2}$$

Option 3:

$$\frac{\pi}{2} - \frac{1}{2}$$

Option 4:

None of these

Correct Answer:

$$\frac{\pi}{4} + \frac{1}{2}$$

Solution:

Simultaneous Trigonometric Equations -

Simultaneous Trigonometric Equations

We can divide the problems related to Simultaneous Trigonometric Equations into two types:

1. If two equations satisfy simultaneously having only one unknown.
2. If two equations satisfy simultaneously having two unknowns.

- _____

$$\sqrt{\cos(1-x)} = \sqrt{\sin x}$$

$$\cos(1-x) \geq 0 \text{ and } \sin x \geq 0$$

$$\cos(1-x) = \sin x$$

$$\sin\left(\frac{\pi}{2} - (1-x)\right) = \sin x$$

$$\frac{\pi}{2} - 1 + x = n\pi + (-1)^n x$$

$$\text{at } n = 1$$

$$2x = \frac{\pi}{2} + 1$$

$$x = \frac{\pi}{4} + \frac{1}{2}$$

For this value of x both satisfies $\cos(1-x) \geq 0$ and $\sin x \geq 0$

Q. 12 The area bounded by $y = \cos x$ and x axis between the ordinates $x = 0$ and $x = \frac{3\pi}{2}$ is

Option 1:

$$4\sqrt{2} - 2$$

Option 2:

$$4\sqrt{2} + 2$$

Option 3:

$$4\sqrt{2} - 1$$

Option 4:

$$4\sqrt{2} + 1$$

Correct Answer:

$$4\sqrt{2} - 2$$

Solution:

As we learnt in


Area along x axis -

Let $y_1 = f_1(x)$ and $y_2 = f_2(x)$ be two curve then area bounded between the


$x = a$ and $x = b$ is

$$\left| \int_a^b \Delta y \, dx \right| = \left| \int_a^b (y_2 - y_1) \, dx \right|$$

- wherein



Where $\Delta y = f_2(x) - f_1(x)$



$$\begin{aligned}
 \text{Area} &= \int_0^{\pi/4} (\cos x - \sin x) dx + \int_{\pi/4}^{5\pi/4} (\sin x - \cos x) dx + \int_{5\pi/4}^{3\pi/2} (\cos x - \sin x) dx \\
 &= [\sin x + \cos x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{5\pi/4} + [\sin x + \cos x]_{5\pi/4}^{3\pi/2} \\
 &= \sqrt{2}^{-1} + [\sqrt{2} + \sqrt{2}] + 1(-1) + \sqrt{2} \\
 &= 4\sqrt{2} - 2
 \end{aligned}$$

Q. 13 If a^{th} term of a $(3a+1)2^{-a}$ is then sum of its infinite term is

Option 1:

10

Option 2:

8

Option 3:

5

Option 4:

0

Correct Answer:

5

Solution:

As we learnt in

Sum of infinite terms of an AGP -

$$S_{\infty} = \frac{a}{1-r} + \frac{dr}{(1-r)^2}$$

here $|r| < 1$

- wherein

$a \rightarrow$ 1st term

$d \rightarrow$ common diff. of a AP

$r \rightarrow$ common ratio of a GP

$$S_{\infty} = \frac{3}{2} + \frac{5}{2^2} + \frac{7}{2^3} + \dots \infty$$

$$\frac{S_{\infty}}{2} = \frac{3}{2^2} + \frac{5}{2^3} + \dots \infty$$

$$\frac{S_{\infty}}{2} = \frac{3}{2} + \frac{1}{2} + \frac{1}{2^2} + \dots \infty$$

$$\frac{S_{\infty}}{2} = \frac{3}{2} + 1$$

$$S_{\infty} = 5$$

Q. 14 Let $\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$ and $\vec{a} + \vec{b} = \hat{i} - \hat{j} - \hat{k}$, then \vec{b} equals

Option 1:

$$\hat{i} + 4\hat{j} - 2\hat{k}$$

Option 2:

$$-\hat{i} - 4\hat{j} - 2\hat{k}$$

Option 3:

$$\hat{i} - 4\hat{j} + 2\hat{k}$$

Option 4:

$$-\hat{i} - 4\hat{j} + 2\hat{k}$$

Correct Answer:

$$-\hat{i} - 4\hat{j} - 2\hat{k}$$

Solution:

G i v e n

$$\vec{a} + \vec{b} = (\hat{i} - \hat{j} - \hat{k})$$

Subtracting vector a from both sides

$$\Rightarrow \vec{b} = (\hat{i} - \hat{j} - \hat{k}) - \vec{a}$$

$$\Rightarrow \vec{b} = (\hat{i} - \hat{j} - \hat{k}) - (2\hat{i} + 3\hat{j} + \hat{k})$$

$$\Rightarrow \vec{b} = -\hat{i} - 4\hat{j} - 2\hat{k}$$

Q. 15

If the fourth term in the Binomial expansion of $\left[\frac{2}{x} + x^{\log_8 x}\right]^6$ is 20×8^7 , then a value of x is :

Option 1:

$$8^3$$

Option 2:

$$8^2$$

Option 3:

$$8$$

Option 4:

$$8^{-2}$$

Correct Answer:

$$8^2$$

Solution:

General Term in the expansion of $(x+a)^n$

$$T_{r+1} = {}^n C_r \cdot x^{n-r} \cdot a^r$$

N o w ,

$$\left[\frac{2}{x} + x^{\log_8 x}\right]^6$$

$$T_4 = T_{3+1} = 20 \times 8^7$$

$$\Rightarrow {}^6C_3 \left(\frac{2}{x}\right)^3 (x^{\log_8 x})^3 = 20 \times 8^7$$

$$\left(\frac{2}{x}\right)^3 (x^{\log_8 x})^3 = (2^3)^7$$

$$\left(\frac{2}{x}\right) (x^{\log_8 x}) = (2)^7$$

$$\frac{x^{\log_8 x}}{x} = 2^6 = 8^2$$

Take log both side with base 8

$$(\log_8 x)^2 = 2 + \log_8 x$$

$$\log_8 x = 2 \text{ or } -1$$

$$x = 8^2$$

Q.16 The vector

$$\left(\hat{i} \times \vec{a} \cdot \vec{b}\right) \hat{i} + \left(\hat{j} \times \vec{a} \cdot \vec{b}\right) \hat{j} + \left(\hat{k} \times \vec{a} \cdot \vec{b}\right) \hat{k} \text{ is equal to :}$$

Option 1:

$$\vec{b} \times \vec{a}$$

Option 2:

$$\vec{A}$$

Option 3:

$$\vec{a} \times \vec{b}$$

Option 4:

$$\vec{B}$$

Correct Answer:

$$\vec{a} \times \vec{b}$$

Solution:

$$\begin{aligned}
& (\hat{i} \times \vec{a} \cdot \vec{b})\hat{i} + (\hat{j} \times \vec{a} \cdot \vec{b})\hat{j} + (\hat{k} \times \vec{a} \cdot \vec{b})\hat{k} \\
& = (\hat{i} \cdot \vec{a} \times \vec{b})\hat{i} + (\hat{j} \cdot \vec{a} \times \vec{b})\hat{j} + (\hat{k} \cdot \vec{a} \times \vec{b})\hat{k} \\
& (\because \vec{a} \times \vec{b} \cdot \vec{c} = \vec{a} \cdot \vec{b} \times \vec{c}) \\
& = (\vec{a} \times \vec{b})\hat{i} + (\vec{a} \times \vec{b})\hat{j} + (\vec{a} \times \vec{b})\hat{k} \\
& = \vec{a} \times \vec{b}
\end{aligned}$$

Q.17 Which of the following inequalities is NOT ne?

Option 1:

$$\int_a^b x \sin x dx \leq \sqrt{\int_a^b x^2 dx \int_a^b \sin^2 x dx}$$

Option 2:

$$\int_a^b \sin x dx \leq \sqrt{\int_a^b \sin^2 x dx}$$

Option 3:

$$\int_a^b x \cos x dx \leq \sqrt{x^3 \int_a^b \cos^2 x dx}$$

Option 4:

$$\int_a^b \ln x \cdot e^x \leq \sqrt{\int_a^b \ln^2 x dx \times \int e^{2x} dx}$$

Correct Answer:

$$\int_a^b x \cos x dx \leq \sqrt{x^3 \int_a^b \cos^2 x dx}$$

Solution:

As we learned

Schwarz - Bunyakovsky Inequality -

If $f^2(x)$ and $g^2(x)$ are integrable in $[a, b]$ then

then

$$\left| \int_a^b f(x) g(x) dx \right| \leq \sqrt{\int_a^b f^2(x) dx \int_a^b g^2(x) dx}$$

-

(c) Violates the Cauchy Schwarz-Bunyakovsky Inequality

Q.18

Let E and F be two independent events. The probability is $\frac{1}{12}$ and the probability that neither E nor F happens is $\frac{1}{2}$

Option 1:

$$\frac{7}{12}$$

Option 2:

$$\frac{3}{2}$$

Option 3:

$$\frac{1}{3}$$

Option 4:

$$\frac{5}{12}$$

Correct Answer:

$$\frac{7}{12}$$

Solution:

Let $P(E) = x$ and $P(F) = y$

$$P(E \cap F) = P(E).P(F) = \frac{1}{12}$$

$$xy = \frac{1}{12}$$

and $P(\text{neither } E \text{ nor } F \text{ happens})$

$$= P(E' \cap F')$$

$$= P(E').P(F')$$

$$= (1 - P(E)).(1 - P(F))$$

$$= (1 - x)(1 - y)$$

Given that $P(\text{neither } E \text{ nor } F \text{ happens}) = 1/2$

$$\text{So } (1 - x)(1 - y) = 1/2$$

$$1 - (x + y) + xy = \frac{1}{2}$$

$$x + y = \frac{7}{12}$$

Q.19 If A is an 3 x 3 non - singular matrix such that $ABA^{-1} = A^{-1}B$ then $B^{-1}A$ is equal to

Option 1:

$$B^{-1}$$

Option 2:

$$(B^{-1})^{-1}$$

Option 3:

$$I + B$$

Option 4:

$$I$$

Correct Answer:

$$I$$

Solution:

As we learnt in

Property of Transpose -

$$(AB)' = B'A'$$

- wherein

A' is the conjugate matrix of A

$$\Rightarrow AA^1 = A^1A$$

$$\text{and } \beta = A^{-1}A^1 \text{ (given)}$$

$$\text{So that } \beta A = A^{-1}A^1A \text{ [multiply by } A]$$

$$= A^{-1}AA^1$$

$$= IA^1$$

$$= A^1$$

$$\text{Now } (BA)^1 = (A^1)^1 = A$$

$$A^1B^1 = A$$

$$A^{-1}A^1B^1 = A^{-1}A = I$$

$$\beta B^1 = I$$

[since $A^{-1}A = I$ and multiply by A^{-1}]

Q. 20 If $A \subseteq B$ and $C \subseteq D$, also $A \times B = P$; $B \times C = Q$ $A \times C = R$; $B \times D = S$. Then

Option 1:

$$P \subseteq S$$

Option 2:

$$Q \subseteq S$$

Option 3:

$$R \subseteq S$$

Option 4:

$$P \subseteq S$$

Correct Answer:

$$R \subseteq S$$

Solution:

As we have learnt,

Theorem of Cartesian Product -

If $A \subseteq B$ and $C \subseteq D$, then $(A \times C) \subseteq (B \times D)$.

-

$$\text{If } A \subseteq B \text{ and } C \subseteq D$$

$$\text{then } A \times C \subseteq B \times D$$

$$\therefore R \subseteq S$$

Q. 21 If $\sec^{-1} \sec(\pi/2 + \theta) = \phi$ where $\theta \in [-\pi/2, \pi/2] - \{0\}$

Then $\phi - \theta = ?$

Option 1:

$$-\pi/2$$

Option 2:

$$\pi/2$$

Option 3:

π

Option 4:

0

Correct Answer:

$\pi/2$

Solution:

As we have learned

Important Results of Inverse Trigonometric Functions -

$$\sec^{-1}(\sec \Theta) = \Theta$$

- wherein

if $0 \leq \Theta < \frac{\pi}{2}$

or $\frac{\pi}{2} < \Theta \leq \pi$

Since $\sec^{-1} \sec \alpha = \alpha$

Thus $\pi/2 + \theta = \phi \Rightarrow \phi - \theta = \pi/2$

Q. 22 A plane passing through $(0, 1, 0)$ and $(0, 0, 1)$ and making $\frac{\pi}{4}$ with the plane $y - z + 5 = 0$, also passes through the point

Option 1:

$(-\sqrt{2}, 1, -4)$

Option 2:

$(\sqrt{2}, -1, 4)$

Option 3:

$(-\sqrt{2}, -1, -4)$

Option 4:

$(\sqrt{2}, 1, 4)$

Correct Answer:

$$(\sqrt{2}, 1, 4)$$

Solution:

Let equation of plane be $a(x - 0) + b(y - 1) + c(z - 0) = 0$

$$\Rightarrow ax + by + cz + b = 0$$

As the given plane also passes through $(0, 0, 1)$

$$c = -b$$

\therefore equation of plane is

$$ax + by - bz + b = 0$$

Also given that this plane makes angle $\frac{\pi}{4}$ with the plane $x + y + z = 0$

$$\cos \frac{\pi}{4} = \frac{a + b + b}{\sqrt{a^2 + b^2 + b^2} \sqrt{2}} = \frac{1}{\sqrt{2}}$$

$$a = \pm \sqrt{2}b$$

Now equation of the plane becomes

$$\pm \sqrt{2}bx + by - bz + b = 0$$

$$\Rightarrow \pm \sqrt{2}x + y - z + 1 = 0$$

$(\sqrt{2}, 1, 4)$ satisfies this equation

Q. 23

The cost of running a bus is Rs. $\left(u + \frac{b}{v}\right)$ per km/h, where u is average speed of the bus. when the bus travels at 30 km/h, the cost comes out to be Rs. 65. Then the most economical speed (in km/h) of the bus is

Option 1:

45

Option 2:

50

Option 3:

60

Option 4:

40

Correct Answer:

60

Solution:

$$\text{Let } \cos \theta = \frac{b}{v}$$

According to the question

$$30a + \frac{b}{30} = 75 \quad \dots (i)$$

$$40a + \frac{b}{40} = 65 \quad \dots (ii)$$

$$a = 1/2 \text{ and } b = 1800$$

$$\text{Now, } C = av + \frac{b}{v}$$

$$\Rightarrow \frac{dC}{dv} = a - \frac{b}{v^2}$$

$$\frac{dC}{dv} = 0 \Rightarrow a - \frac{b}{v^2} = 0$$

$$\Rightarrow v = \sqrt{\frac{b}{a}} = \sqrt{3600}$$

$$\Rightarrow y = 60 \text{ kmph}$$

Q. 24 Find $\int \frac{dx}{x^2 - 4}$

Option 1:

$$\ln \frac{x+2}{x-2} + C$$

Option 2:

$$1/2 \ln \frac{x-2}{x+2} + C$$

Option 3:

$$\frac{1}{4} \ln \left| \frac{x-2}{x+2} \right| + C$$

Option 4:

none of these

Correct Answer:

$$\frac{1}{4} \ln \left| \frac{x-2}{x+2} \right| + C$$

Solution:

As we have learned

Rule of integration by Partial fraction -

Linear and non-repeated:

$$\frac{P(x)}{Q(x)} = \frac{P(x)}{(x - \alpha_1)(x - \alpha_2) \cdots (x - \alpha_n)}$$

$$\text{Let } \frac{P(x)}{Q(x)} = \frac{A}{(x - \alpha_1)} + \frac{B}{(x - \alpha_2)} \cdots$$

Find a, b, \dots

By comparing $\frac{P(x)}{Q(x)}$

-

$$\begin{aligned} I &= \int \frac{dx}{x^2 - 4} = \int \frac{dx}{(x - 2)(x + 2)} \\ &= -\frac{\ln(|x + 2|) - \ln(|x - 2|)}{4} + C \\ &= \frac{1}{4} \ln \left| \frac{x - 2}{x + 2} \right| + C \end{aligned}$$

Q. 25

$$\text{If } S = \left\{ x \in [0, 2\pi] : \begin{vmatrix} 0 & \cos x & -\sin x \\ \sin x & 0 & \cos x \\ \cos x & \sin x & 0 \end{vmatrix} = 0 \right\},$$

then $\sum_{x \in S} \tan\left(\frac{\pi}{3} + x\right)$ is equal to :

Option 1:

$$4 + 2\sqrt{3}$$

Option 2:

$$-2 + \sqrt{3}$$

Option 3:

$$-2 - \sqrt{3}$$

Option 4:

$$-4 - 2\sqrt{3}$$

Correct Answer:

$$-2 - \sqrt{3}$$

Solution:

As we learnt in

Cramer's rule for solving system of linear equations -

When $\Delta = 0$ and $\Delta_1 = \Delta_2 = \Delta_3 = 0$,

then the system of equations has infinite solutions.

- wherein

$$a_1x + b_1y + c_1z = d_1$$

$$a_2x + b_2y + c_2z = d_2$$

$$a_3x + b_3y + c_3z = d_3$$

and

$$\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$\Delta_1, \Delta_2, \Delta_3$ are obtained by replacing (d_1, d_2, d_3) in each column of

$$\Rightarrow \begin{vmatrix} 0 & \cos x & -\sin x \\ \sin x & 0 & \cos x \\ \cos x & \sin x & 0 \end{vmatrix} = 0$$

$$\Rightarrow 0(0 - \sin x \cos x) - \cos x(0 - \cos^2 x) - \sin x(\sin^2 x) = 0$$

$$\Rightarrow \sin^3 x - \cos^3 x = 0$$

$$\Rightarrow (\sin x - \cos x)(1 + \sin x \cos x) = 0$$

$$\Rightarrow \therefore \sin x - \cos x = 0$$

$$\Rightarrow \therefore \sin x - \cos x = 0$$

$$\therefore \tan x = 1$$

$$\therefore x = \frac{\pi}{4} \text{ or } \frac{5\pi}{4}$$

$$\text{Now } \tan\left(\frac{\pi}{4} + x\right) = \frac{1 + \tan x}{1 - \tan x}$$

$$\text{Now put } \frac{\pi}{3}$$

$$= \frac{1 + \tan \frac{\pi}{3}}{1 - \tan \frac{\pi}{3}}$$

$$= \frac{1 + \sqrt{3}}{1 - \sqrt{3}}$$

$$= \frac{(1 + \sqrt{3})^2}{1 - 3} = \frac{1 + 3 + 2\sqrt{3}}{-2} = -2 - \sqrt{3}$$

$$\sum \tan\left(\frac{\pi}{3} + x\right) = -2 - \sqrt{3}$$

Q.26 Let W denote the words in the English Dictionary. Define a relation R on W as follows:

$R = \{(x, y) \in W \times W \mid \text{the word } x \text{ and } y \text{ have at least one letter in common}\}$. Then R is

Option 1:

not reflexive, symmetric and transitive

Option 2:

reflexive, symmetric and not transitive

Option 3:

reflexive, symmetric and transitive

Option 4:

reflexive, not symmetric and transitive.

Correct Answer:

reflexive, symmetric and not transitive

Solution:

As we learnt in

REFLEXIVE RELATION: A relation R in A is said to be reflexive, if $(a, a) \in R$ for every $a \in A$.

SYMMETRIC RELATION: A relation R in A is said to be symmetric, if $(a, b) \in R$ implies $(b, a) \in R$ for every $a, b \in A$.

TRANSITIVE RELATION: A relation R in A is said to be transitive, if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for every $a, b, c \in A$.

Now,

$$R = \{(x, y) \in \omega \times \omega\}$$

x, y having at least one letter in common

1. Reflexive

For any pair of type $(\text{word } 1, \text{word } 1)$, both are same, so they lie in the relation. So, the relation is reflexive.

2. Symmetric

If $(\text{word } 1, \text{word } 2)$ lie in this relation, then $(\text{word } 2, \text{word } 1)$ also lies in the relation, the relation is symmetric

If $(\text{word } 1, \text{word } 2)$ lies in R , then word 1 and word 2 have at least one letter in common

So, definitely word 2 and word 1 have at least one letter in common relation

So, if $(\text{word } 1, \text{word } 2)$ lies in R , then $(\text{word } 2, \text{word } 1)$ also lies in R

So, it is a symmetric relation

3. Transitive

If $(\text{word } 1, \text{word } 2)$ lies in R , then they have at least one letter in common

If $(\text{word } 2, \text{word } 3)$ lies in R , so these 2 words have at least one letter in common

But, this does not mean that $(\text{word } 1, \text{word } 3)$ lies in R , as they may not have any letter in common. For example, (cat, bat) , and $(\text{bat}, \text{blur})$ lie in R , but $(\text{cat}, \text{blur})$ does not lie in R .

So, it is not transitive.

Correct option is 2.

Q.27 The set of all real values of λ for which exactly two common tangents exist to the circles $x^2 + y^2 - 4x - 4y + 6 = 0$ and $x^2 + y^2 - 10x - 10y + \lambda = 0$

Option 1:

(1 2 , 3 2)

Option 2:

(1 8 , 4 2)

Option 3:

(1 2 , 2 4)

Option 4:

(1 8 , 4 8)

Correct Answer:

(1 8 , 4 2)

Solution:

As we learnt, when 2 circles intersect at 2 distinct points, then there are two common tangents, both of them being direct.

The condition for this is $|r_1 - r_2| < C_1C_2 < r_1 + r_2$

N o w ,

$$S_1 : x^2 + y^2 - 4x - 4y + 6 = 0$$

$$r_1 = \sqrt{2^2 + 2^2 - 6} = \sqrt{2}$$


$$C_1 : (2, 2)$$

$$S_2 : x^2 + y^2 - 10x - 10y + \lambda = 0$$

$$r_2 = \sqrt{5^2 + 5^2 - \lambda} = \sqrt{50 - \lambda}$$

$$C_2 : (5, 5)$$

F o r t w o t a n g e n t s



$$|r_1 - r_2| < C_1C_2 < r_1 + r_2$$

$$|\sqrt{50 - \lambda} - \sqrt{2}| < 3\sqrt{2} < \sqrt{2} + \sqrt{50 - \lambda}$$

$$\text{W e } \sqrt{50 - \lambda} < 4\sqrt{2} \Rightarrow \lambda > 18 \text{ and } \lambda \leq 50$$

$$\sqrt{50 - \lambda} > 2\sqrt{2} \Rightarrow \lambda < 42$$

$$\lambda \in (18, 42)$$

Q. 28 If $f(x) = 3|x|$. Then the range of $f(x)$ is

Option 1:

$[0, \infty)$

Option 2:

$[3, \infty)$

Option 3:

\mathbb{R}

Option 4:

$[1, \infty)$

Correct Answer:

$[0, \infty)$

Solution:

We can find range of $|x|$ by simple manipulation

$$As \quad 0 \leq |x| < \infty$$

Multiplying all 3 sides by 3

$$0 \leq 3|x| < \infty$$

So, range is the same i.e $[0, \infty)$

Q. 29 What is the union of set $A = \{1, 2, 3, 4\}$ with a set $B = \{x : x^2 - 5x + 6 = 0 \text{ or } x^2 - 5x + 4 = 0\}$

Option 1:

A

Option 2:

B

Option 3:

Both (1) and (2)

Option 4:

None of these

Correct Answer:

Both (1) and (2)

Solution:

As we learnt in

UNION OF SETS -

Let A and B be any two sets. The union of A and B is the set with the elements of A and B, the common elements being taken only once.

And $A \cup A = A$

In this Question,

$$B = \{x : x^2 - 5x + 6 = 0 \text{ or } x^2 - 5x + 4 = 0\}$$

$$x^2 - 5x + 6 = 0 \Rightarrow x = 2, 3$$

$$x^2 - 5x + 4 = 0 \Rightarrow x = 1, 4$$

$$\Rightarrow B = \{1, 2, 3, 4\}$$

Thus $A = B$

Thus $A \cup B = A = B$

Q. 30 If $A \times B = P; C \times D = Q, A \cap C = R, B \cap D = S$, then which of the following is true ?

Option 1:

$$P \cup Q = R \times S$$

Option 2:

$$P \cap Q = R \times S$$

Option 3:

$$P \times Q = R \cup S$$

Option 4:

$$P \times Q = R \cap S$$

Correct Answer:

$$P \cap Q = R \times S$$

Solution:

As we have learnt,

Theorem of Cartesian Product -

$$(A \times B) \cap (C \times D) = (A \cap C) \times (B \cap D)$$

-

$$(A \times B) \cap (C \times D) = (A \cap C) \times (B \cap D)$$
$$P \cap Q = R \times S$$

Q. 31 A circular wire of radius 3 cm is cut and bent so as to l
the angle subtended by the wire at the center of the ci

Option 1:

$$\frac{\pi}{8}$$

Option 2:

$$\frac{\pi}{4}$$

Option 3:

$$\frac{\pi}{6}$$

Option 4:

None of these

Correct Answer:

$$\frac{\pi}{8}$$

Solution:

Length of the circular wire = $2\pi r = 6\pi$ cm.

Angle subtended by the arc at the centre = $\frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \frac{6\pi}{48} = \frac{\pi}{8} \text{ radian}$$

Q. 32 If the truth value of the statement $p \leftrightarrow (p \vee q \vee r)$ is false (F), then the truth v
statements p, q, r are respectively :

Option 1:

F, T, T

Option 2:

T, F, F

Option 3:

T, T, F

Option 4:

T, F, T

Correct Answer:

T, T, F

Solution:

Tautology And Contradiction -

Tautology

A compound statement is called a tautology if it is always true for all possible truth values of its component statements.

For example, $(p \Rightarrow q) \vee (q \Rightarrow p)$

Contradiction (fallacy)

A compound statement is called a contradiction if it is always false for all possible truth values of its component statements.

For example, $(p \Rightarrow q) \vee (q \Rightarrow p)$

Truth Table

p	q	$p \rightarrow q$	$q \rightarrow p$	$(p \rightarrow q) \vee (q \rightarrow p)$	$\sim ((p \rightarrow q) \vee (q \rightarrow p))$
T	T	T	T	T	F
T	F	F	T	T	F
F	T	T	F	T	F
F	F	T	T	T	F

-

$$p \rightarrow (\sim q \vee r)$$

$$\sim p \vee (\sim q \vee r)$$

$$\left. \begin{array}{l} \sim p \rightarrow F \\ \sim q \rightarrow F \\ r \rightarrow T \end{array} \right\} \Rightarrow \begin{array}{l} p \rightarrow T \\ q \rightarrow T \\ r \rightarrow F \end{array}$$

Q. 33 A variable circle passes through the fixed point $A(p, q)$ and touches the x -axis. The locus of the other end of the diameter through A is

Option 1:

$$(y - p)^2 = 4qx$$

Option 2:

$$(x - q)^2 = 4py$$

Option 3:

$$(x - p)^2 = 4qy$$

Option 4:

$$(y - q)^2 = 4px$$

Correct Answer:

$$(x - p)^2 = 4qy$$

Solution:

As we learnt in

Equation of a circle -

$$(x - h)^2 + (y - k)^2 = r^2$$

- wherein

Circle with centre (h, k) and radius r .

Circle touching x -axis and having radius r -

$$x^2 + y^2 \pm 2rx + 2fy + f^2 = 0$$

- wherein

Where f is a variable parameter.

Let the other diametric end be $P(h, k)$

So centre is $\left(\frac{p+h}{2}, \frac{q+R}{2}\right)$

Radius $= \sqrt{\left(\frac{h-p}{2}\right)^2 + \left(\frac{k-q}{2}\right)^2}$

For circle touching y -axis, radius = $\frac{q+k}{2}$

$$\text{So } \left(\frac{h-p}{2}\right)^2 + \left(\frac{k-q}{2}\right)^2 = \left(\frac{k+q}{2}\right)^2$$

we get $(h-p)^2 = 4kq$

i.e. $(x-p)^2 = 4qy$. a parabola

Q. 34 If $f(x) = \log_e \left(\frac{1-x}{1+x} \right)$, $|x| < 1$, then $f\left(\frac{2x}{1+x^2}\right)$ is equal to:

Option 1:

$2f(x^2)$

Option 2:

$2f(x)$

Option 3:

$-2f(x)$

Option 4:

$(f(x))^2$

Correct Answer:

$2f(x)$

Solution:

$$f(x) = \log_e \left(\frac{1-x}{1+x} \right), |x| < 1,$$

$$f\left(\frac{2x}{1+x^2}\right) = ?$$

$$f\left(\frac{2x}{1+x^2}\right) = \log_e \left(\frac{1 - \frac{2x}{1+x^2}}{1 + \frac{2x}{1+x^2}} \right)$$

$$\begin{aligned}
&= \log_e \left(\frac{\frac{1+x^2-2x}{1+x^2}}{\frac{1+x^2+2x}{1+x^2}} \right) \\
&= \log_e \frac{x^2 - 2x + 1}{x^2 + 2x + 1} = \log_e \left(\frac{(1-x)^2}{(x+1)^2} \right) \\
&= \log_e \left(\frac{1-x}{x+1} \right)^2 \\
&= 2 \log_e \left(\frac{1-x}{x+1} \right), \quad |x| < 1 \\
&= 2f(x)
\end{aligned}$$

Q. 35 If $A \times B = \{1, 3, 4\}$ and $A \times C = \{3, 4\}$ then $A \times (C - B) =$

Option 1:

$\{1\}$

Option 2:

$\{3, 4\}$

Option 3:

$\{1, 3, 4\}$

Option 4:

None of these.

Correct Answer:

None of these.

Solution:

As we have learnt,

Theorem of Cartesian Product -

$$A \times (B - C) = (A \times B) - (A \times C)$$

-

$$A \times (C - B) = (A \times C) - (A \times B)$$

= doesn't exist =

Q. 36 Solution of differential equation is

Option 1:

$$e^{x^2/2}(\sin y - 1) = c$$

Option 2:

$$e^x(\sin y - 1) = c$$

Option 3:

$$e^{x^2}(\sin y - 1) = c$$

Option 4:

$$e^{x^2}(\sin y + 1) = c$$

Correct Answer:

$$e^{x^2/2}(\sin y - 1) = c$$

Solution:

As we have learned

Extended Form of linear Differential Equation -

Sometimes, a differential equation is not linear but it can be -

$$\text{Let } \sin y = t \Rightarrow \cos y \frac{dy}{dx} = \frac{dt}{dx}$$

equation reduces to

$$\frac{dt}{dx} + (x)t = x$$

which is again comparable with

$$I.F = e^{\int x dx} = e^{x^2/2}$$

multiplying both sides with integrating factor, we get

$$e^{x^2/2} \cdot \frac{dt}{dx} + t e^{x^2/2} \cdot x = x e^{x^2/2}$$

$$\Rightarrow d/dx(e^{x^2/2} \cdot t) = x e^{x^2/2}$$

$$\Rightarrow \int d/dx(e^{x^2/2} \cdot t) - \int x e^{x^2/2} dx = C$$

$$\Rightarrow e^{x^2/2} \cdot t - e^{x^2/2} = c$$

$$\Rightarrow e^{x^2/2} \cdot (t - 1) = C \Rightarrow e^{x^2/2}(\sin y - 1) = C$$

Q. 37 If the tangent to the curve, the points perpendicular to the line, $x + y + 4 = 0$, then which one of the following points lies

Option 1:

$(-2, 1)$

Option 2:

$(-2, 2)$

Option 3:

$(2, -1)$

Option 4:

$(2, -2)$

Correct Answer:

$(2, -2)$


Solution:

Slope of the tangent -

Let $y = f(x)$ is a curve then $dy / dx = f'(x)$ and at a particular

$$M_T = \lim_{\delta x \rightarrow 0} \frac{(y + \delta y) - y}{(x + \delta x) - x} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x}$$

- wherein



Equation of tangent -

Equation of tangent from other point on the curve.

Let $y = mx + c$ and (h, k) is the point where it passes through itself. Then
 $C = K - mh$

$$y = mx + K - mh$$

- wherein

Where m is slope of tangent.

given,

$$y = x^3 + ax - b$$

If pass (1, -5) i.e.

$$\Rightarrow -5 = 1 + a - b \Rightarrow a - b = -6. \dots \dots \dots (1)$$

Slope of the tangent = 0 i.e.

Slope of the tangent = 0 i.e.

$$\frac{dy}{dx} = 3x^2 + a$$

$$\Rightarrow 3 + a = -1 \quad x = 1 \quad \text{at}$$

$$\Rightarrow a = -4. \dots \dots \dots (2)$$

from (1) and (2)

$$a = -4, b = +2$$

equation of curve become

$$y = x^3 - 4x - 2$$

$(2, -2)$ satisfies the equation

Q. 38 Let $\vec{a}, \vec{b}, \vec{c}, \vec{d}$ are four vectors then $[\vec{a} + \vec{b} \ \vec{c} \ \vec{d}] + [\vec{b} + \vec{c} \ \vec{a} \ \vec{d}] + [\vec{a} + \vec{c} \ \vec{b} \ \vec{d}]$ equals

Option 1:

- 1

Option 2:

0

Option 3:

1

Option 4:

2

Correct Answer:

0

Solution:

As we have learned

Properties of Scalar Triple Product -

$$\left[(\vec{a} + \vec{b}) \vec{c} \vec{d} \right] = \left[\vec{a} \vec{c} \vec{d} \right] + \left[\vec{b} \vec{c} \vec{d} \right]$$

- wherein

$\vec{a}, \vec{b}, \vec{c}, \vec{d}$ are four vectors.

$$\begin{aligned} & \left[\vec{a} + \vec{b} \vec{c} \vec{d} \right] + \left[\vec{b} + \vec{c} \vec{a} \vec{d} \right] + \left[\vec{a} + \vec{c} \vec{b} \vec{d} \right] \\ &= \left[\vec{a} \vec{c} \vec{d} \right] + \left[\vec{b} \vec{c} \vec{d} \right] + \left[\vec{b} \vec{a} \vec{d} \right] + \left[\vec{c} \vec{a} \vec{d} \right] + \left[\vec{a} \vec{b} \vec{d} \right] + \left[\vec{c} \vec{b} \vec{d} \right] \\ &= \left[\vec{a} \vec{c} \vec{d} \right] + \left[\vec{b} \vec{c} \vec{d} \right] + \left[\vec{b} \vec{a} \vec{d} \right] - \left[\vec{a} \vec{c} \vec{d} \right] - \left[\vec{b} \vec{a} \vec{d} \right] - \left[\vec{b} \vec{c} \vec{d} \right] = 0 \end{aligned}$$

Q. 39 Which of the following integral has maximum value?

Option 1:

$$\int_0^5 \ln x dx$$

Option 2:

$$\int_0^5 |\ln x| dx$$

Option 3:

both (a) and (b) are equal

Option 4:

$$5 \int_0^1 \ln x dx$$

Correct Answer:

$$\int_0^5 |\ln x| dx$$

Solution:

As we have learnt,

Properties of Definite Integration -

If $f(x)$ is defined on (a, b) then $\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$

-

Graph of $|\ln x|$ is always above the x-axis.

So it encloses the maximum area under the curve.

Q.40 The volume of the largest possible right circular cylinder of radius $\sqrt{3}$ is :

Option 1:

$$\frac{4}{3}\sqrt{3}\pi$$

Option 2:

$$\frac{8}{3}\sqrt{3}\pi$$

Option 3:

$$4\pi$$

Option 4:

$$2\pi$$

Correct Answer:

$$4\pi$$

Solution:

As we have learned

Method for maxima or minima -

By second derivative method :

Step 1. find values of x for $\frac{dy}{dx} = 0$

Step 2. $x = x_0$ is a point of local maximum if $f''(x) < 0$ and local minimum if $f''(x) > 0$

- wherein


Where $y = f(x)$

$$\frac{dy}{dx} = f'(x)$$

vol. of cylinder

$$\text{Now, } R^2 = r^2 + \frac{h^2}{4}$$

$$\Rightarrow 3 = r^2 + \frac{h^2}{4}$$



$$\Rightarrow h^2 = 4(3 - r^2)$$
$$\Rightarrow r^2 = 3 - (h^2/4)$$

$$V = \pi h \left(3 - \frac{h^2}{4} \right)$$

$$= 3\pi h - \frac{\pi}{4} h^3$$

$$\Rightarrow \frac{dv}{dh} = 3\pi - \frac{3\pi}{4} h^2$$

$$\Rightarrow h = 2$$

$$\therefore \text{volume} = \pi \times 2 \times (3 - 1) = 4\pi$$

English

Q. 1 Find out the meaning of the phrasal verb

Break in

Option 1:

Enter like a thief

Option 2:

Ask one to do something

Option 3:

Interrupt

Option 4:

Trouble

Option 5:

Ask for a favour

Correct Answer:

Interrupt

Solution:

The verb means to interrupt

Q. 2 Read the paragraph and choose a suitable summary from the given options:

Journalism as a profession is gaining ground in these days. There is an explosion of information. We are living in a fast changing world with a fast flow of information. A large number of magazines and journals have come up. Today, we have a number of news agencies trading in news and there are more than two scores of associations of journalists in the country. There are regular courses in journalism run in various universities. Mass communication has already been introduced as a subject at post graduate level in many universities. The profession of journalism now attracts a lot of able and talented students.

Option 1:

Journalism has always been a rampantly growing force

Option 2:

Journalism as a calling is going through a period of considerable snowballing

Option 3:

Journalism is the best profession for students now a days.

Option 4:

Journalism is unbridled and thus, perilous

Option 5:

Journalism is defying the growth of all other professions

Correct Answer:

Journalism as a calling is going through a period of considerable snowballing

Solution:

The paragraph highlights the expansion and growth of journalism. "Journalism as a calling is going through a period of considerable snowballing" is the best choice.

Q.3 Change the speech.

She said, "I am sorry, I did not write".

Option 1:

She apologized for I did not write.

Option 2:

She apologized she is sorry, she could not write.

Option 3:

She apologized and added that she was sorry, she had not written.

Option 4:

She apologized she is sorry, she did not write.

Option 5:

She apologized she is sorry, she has not written.

Correct Answer:

She apologized and added that she was sorry, she had not written.

Solution:

grammatically and idiomatically correct option is 4.

Q.4

Find out the pair of words from the options which exhibit the same relationship as the pair of words:

Lull: Break

Option 1:

Destitution: Penury

Option 2:

Paucity: Abundance

Option 3:

Agent: Penurious

Option 4:

Misogamist: Brawn

Correct Answer:

Destitution: Penury

Solution:

Lull means a break. Similarly, destitution means penury.

Q.5

Read the paragraph and choose a suitable summary from the given options: Does any innocent

human being deserve to be abused? The answer is no. So why should animals on factory farms have to go through this pain? They feel and hurt too, as they're living things. That's why people should focus on animal protection on factory farms because the treatment is cruel and immoral. First of all, animals suffer everywhere because of what people do to them. So hidden cameras are set up in all different factory farms all over to further prove the wrong that's happening.

Option 1:

Guiltless animals are subjected to inhumane behaviour by human beings and thus, they need to be resuscitated.

Option 2:

Animals are innocuous and thus, become the target of savagery at the hands of human beings.

Option 3:

Killing animals is inhumane and thus, must be discouraged.

Option 4:

Innocent animals are killed and it shows inhuman nature of

Option 5:

It is imperative to save animals to maintain the equilibrium of the eco system.

Correct Answer:

Animals are innocuous and thus, become the target of savagery at the hands of human beings.

Solution:

The author underlines that animals are innocent, brutalized and a comprehensive choice is “Animals are innocuous and thus, become the target of savagery at the hands of human beings”

Q. 6 Change the voice:

This is being done by him

Option 1:

This is being done by him

Option 2:

He is doing it

Option 3:

He will do it

Option 4:

He has been doing this

Option 5:

He has done it

Correct Answer:

He is doing it

Solution:

“He is doing it” appropriately changes the subject to object.

Q.7 Change the speech.

Bala's mother, "Will you lock the door, Bala ?"

Option 1:

Bala's mother asked Bala if he would lock the door.

Option 2:

Bala's mother asked Bala if he can lock the door.

Option 3:

Bala's mother said lock the door.

Option 4:

Bala's mother told Bala that he need not lock the door.

Option 5:

Bala's mother asked Bala if he had locked the door.

Correct Answer:

Bala's mother asked Bala if he would lock the door.

Solution:

Rest of the options change the meaning.

Q.8 Find out the correct form of tense.

Employees who fought the adverse conditions narrated control after forty minutes.

Option 1:

was brought

Option 2:

has been brought

Option 3:

is brought

Option 4:

had been brought

Option 5:

can be brought

Correct Answer:

was brought

Solution:

The sentence is in past tense. Was brought thus, is idiomatic

Q. 9 Find out the odd one out among the following words:

Option 1:

Extortionate

Option 2:

Extravagant

Option 3:

Reasonable

Option 4:

Exorbitant

Option 5:

Dear

Correct Answer:

Reasonable

Solution:

Reasonable is the odd word. Where the others denote a high cost or are cheaper.

Q. 10 Find out the synonym for the given word:

Discuss

Option 1:

Sidestep

Option 2:

Enlist

Option 3:

Broach

Option 4:

ill-disposed

Option 5:
Loath

Correct Answer:
Broach

Solution:

Broach means to raise a topic for discussion.

Logical Reasoning

Q.1 Choose the word which is least like the other words in the group

Option 1:
37

Option 2:
45

Option 3:
49


Option 4:
65

Correct Answer:
49


Solution:

49 is the only Perfect square number in the group


Q.2




Option 1:




Option 2:




Option 3:



Option 4:



Correct Answer:




Solution:

The arrow heads pointed towards the center

Q.3

Study the following table carefully to answer the question.
Character.



Option 1:

4

Option 2:

3 0 5

Option 3:

3 4 3

Option 4:

7 2 9


Correct Answer:

3 4 3

Solution:

Moving clockwise, the terms are :

Q.4 How the gure wiil look after 180 degree rotation



Option 1:



Option 2:



Option 3:



Option 4:



Correct Answer:



Solution:

After 180 degree the gure look like

Q.5

Study the following table carefully to answer the question.
Character

594	198
?	66

Option 1:

1 1

Option 2:

1 2

Option 3:

2 2

Option 4:

3 3

Correct Answer:

2 2

Solution:

Moving clockwise we have : $594/3=198$; $198/3=66$;
 $66/3=22$

Q.6

In each question below is given a passage followed by example each inference separately in the context of the degree of truth or falsity.

Corel draw and photoshop are the old softwares used for Adobe illustrator and the In Design are the new ones, but are mostly used in the industries. These are designing for Digital Marketing. Digital Marketing is a technique used for facebook marketing, instagram and E-mail marketing etc. approx 70 % of companies to generate lead for their business end of traditional marketing strategies. By the coming scene of marketing is dramatically changed. In india on internet.

70 % of companies used digital marketing technique to generate their business leads.

Option 1:

if the inference is "definitely true" it directly follows from

Option 2:

if the inference is "probably true" though not definitely t

Option 3:

if you think the data is inadequate, from the facts given you likely to be true or false

Option 4:

if you think the inference is "probably false" though not given

Correct Answer:


if the inference is "probably true" though not definitely t

Solution:

The information is mentioned in the passage.

Q.7

Study the following table carefully to answer the question.
Character.



Option 1:

4 9

Option 2:

7 6

Option 3:

8 9

Option 4:

9 4

Correct Answer:

4 9

Solution:

we have : $(18 + 19) - 2 = 35$;

$(22 + 24) - 3 = 43$

so the answer is : $(26 + 27) - 4 = 49$

Q.8

Bird : Cage

Option 1:

Animals : Zoo

Option 2:

Thief : Prison

Option 3:

Antique : Museum

Option 4:

Crime : Punishment

Correct Answer:
Thief : Prison

Solution:
Second is the place where the first is made captive

Q.9 Train : Track

Option 1:
Idea : Brain

Option 2:
Bullet : Barrel


Option 3:
Water : Boat

Option 4:
Fame : Television


Correct Answer:
Bullet : Barrel

Solution:
A Train moves on tracks. Similarly , a bullet travels through the barrel

Q. 10




Option 1:



Option 2:



Option 3:



Option 4:



Correct Answer:



Solution:

The triangles are increasing by 1

Q.11 In each question below is given a passage followed by example each inference separately in the context of the degree of truth or falsity.

Now, In India the strength of girls in the colleges are increasing because of the several movements led by the institutions. There is another movement which is increasing members. Only 30% of females are doing job and they are government has been opening so many courses to skill them. There is a sharp increment in the ratio of new born girls after the campaign "beti bachao beti padhao" led by our

In India sex ratio is good in comparison to other countries.

Option 1:

if the inference is "probably true" though not definitely true

Option 2:

if the inference is "definitely true" it directly follows from

Option 3:

if you think the data is inadequate , from the facts given you are likely to be true or false

Option 4:

if you think the inference is " probably false " though not definitely false given

Correct Answer:

if you think the data is inadequate , from the facts given you are likely to be true or false

Solution:

The details about the sex ratio of other countries is not discussed.

Q.12 In each question below is given a passage followed by four statements. Choose the statement which is true or false. Example each inference separately in the context of the passage and the degree of truth or falsity.

Corel draw and photoshop are the old softwares used for designing. Adobe illustrator and the In Design are the new ones, but they are mostly used in the industries. These are designing softwares for Digital Marketing. Digital Marketing is a technique used for facebook marketing, instagram and E-mail marketing etc. About 70 % of companies use digital marketing to generate lead for their business. It is the end of traditional marketing strategies. By the coming of digital marketing scene of marketing is dramatically changed. In india online marketing is internet.

Internet has been used in india from year 2000.

Option 1:

if the inference is " probably true " though not definitely true

Option 2:

if the inference is " definitely true " it directly follows from the passage

Option 3:

if you think the data is inadequate , from the facts given you are likely to be true or false

Option 4:

if you think the inference is 'definitely false' it contradicts the passage


Correct Answer:

if you think the inference is 'definitely false' it contradicts the passage

Solution:

In the passage it is mention that internet in india starts from

Q.13 Chart Logic (Insert the Missing Character)



Option 1:



Option 2:



Option 3:



Option 4:




Correct Answer:



Solution:

The arrow pointed towards the center

Q.14 Find out which of the figure formed after combining the



Option 1:



Option 2:



Option 3:



Option 4:



Correct Answer:



Solution:

it is clear from the figure

Q.15 Choose the word which is least like the other words in the

Option 1:

Calender

Option 2:

Year

Option 3:

D a t e

Option 4:

M o n t h

Correct Answer:

C a l e n d e r

Solution:

A l l O t h e r s a r e p a r t s o f a C a l e n d e r

Q. 16 1 , 9 , 2 5 , 4 9 , ? , 1 2 1

Option 1:

6 4

Option 2:

8 1

Option 3:

9 1

Option 4:

1 0 0

Correct Answer:

8 1

Solution:

T h e g i v e n s e r i e s c o n s i s t s o f s q u a r e s o f c o n s e c u t i v e o d d n u m b

$1^2, 3^2, 5^2, \dots$

S o , m i s s i n g ² t e r m =

Q. 17 2 , 3 , 8 , 2 7 , 1 1 2 , ?

Option 1:

2 2 6

Option 2:

3 3 9

Option 3:

4 5 2

Option 4:

5 6 5

Correct Answer:

5 6 5

Solution:

The Pattern is $X_1 + 1$, $X_2 + 2$, $X_3 + 3$, $X_4 + 4$,
so missing term = $112 \times 5 + 5 = 565$

Q.18 Choose the word which is least like the other words in the

Option 1:

Curd

Option 2:

Butter

Option 3:

Oil

Option 4:

Cheese

Correct Answer:

Oil

Solution:

All except Oil are products obtained from milk.

Q.19 Zinc is to Galvanisation as Nickel is to

Option 1:

Aircraft

Option 2:

Corrosion

Option 3:

Electroplating

Option 4:

Filament

Correct Answer:
Electroplating

Solution:

Second is the purpose of rst

Q.20 25 : 125 : : 36 : ?

Option 1:
180

Option 2:
206

Option 3:
216

Option 4:
318

Correct Answer:
216

Solution:

The relationship is $x^2 : x^3$ is